

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 October 2002 (31.10.2002)


PCT

(10) International Publication Number
WO 02/085357 A1


- (51) International Patent Classification⁷: **A61K 31/44**, 31/48, 31/415, 31/445, C07D 211/04, 211/06, 235/24, 401/04, 401/06, 403/04, 403/06, 471/02, 498/02
- (71) Applicant (for all designated States except US): **EURO-CELTIQUE, S.A.** [LU/LU]; 122 Boulevard de la Petrusse, L-2330 Luxembourg (LU).
- (21) International Application Number: **PCT/US02/12351**
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **SUN, Qun** [CN/US]; 68 Hills Drive, Belle Mead, NJ 08502 (US). **GOEHRING, R., Richard** [US/US]; 6195 Potters Lane, Pipersville, PA 18947 (US). **KYLE, Donald** [US/US]; 29 Weatherfield Drive, Newtown, PA 18940 (US). **CHEN, Zhengming** [CN/US]; 11 Hudnut Lane, Belle Mead, NJ 08502 (US). **VICTORY, Sam** [US/US]; 55 Quince Circle, Newtown, PA 18940 (US). **WHITEHEAD, John** [GB/US]; 155 North State Street, Newtown, PA 18940 (US).
- (22) International Filing Date: **18 April 2002 (18.04.2002)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
- | | | |
|------------|----------------------------|----|
| 60/284,666 | 18 April 2001 (18.04.2001) | US |
| 60/284,667 | 18 April 2001 (18.04.2001) | US |
| 60/284,668 | 18 April 2001 (18.04.2001) | US |
| 60/284,669 | 18 April 2001 (18.04.2001) | US |
- (74) Agents: **DAVIDSON, Clifford, M. et al.**; Davidson, Davidson & Kappel, LLC, 485 Seventh Avenue, 14th Floor, New York, NY 10018 (US).

[Continued on next page]

(54) Title: **NOCICEPTIN ANALOGS**




(I)




(II)

(57) Abstract: A compound of formula (I), (II), (III) or (IV), wherein Z, A, B, C, R, R₁, R₂, Q, and n are as described herein.



(III)



(IV)

WO 02/085357 A1



(81) **Designated States (national):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(84) **Designated States (regional):** ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

NOCICEPTIN ANALOGS

This application claims priority from U.S. Provisional Application Serial Nos. 60/284,666; 60/284,667; 60/284,668; 60/284,669 all filed April 18, 2001, the disclosures of which are hereby incorporated by reference.

BACKGROUND OF THE INVENTION

Chronic pain is a major contributor to disability and is the cause of an untold amount of suffering. The successful treatment of severe and chronic pain is a primary goal of the physician with opioid analgesics being preferred drugs.

Until recently, there was evidence of three major classes of opioid receptors in the central nervous system (CNS), with each class having subtype receptors. These receptor classes were designated as μ , δ and κ . As opiates had a high affinity to these receptors while not being endogenous to the body, research followed in order to identify and isolate the endogenous ligands to these receptors. These ligands were identified as enkephalins, endorphins and dynorphins.

Recent experimentation has led to the identification of a cDNA encoding an opioid receptor-like (ORL1) receptor with a high degree of homology to the known receptor classes. This newly discovered receptor was classified as an opioid receptor based only on structural grounds, as the receptor did not exhibit pharmacological homology. It was initially demonstrated that non-selective ligands having a high affinity for μ , δ and κ receptors had low affinity for the ORL1. This characteristic, along with the fact that an endogenous ligand had not yet been discovered, led to the term "orphan receptor".

Subsequent research led to the isolation and structure of the endogenous ligand of the ORL1 receptor. This ligand is a seventeen amino acid peptide structurally similar to members of the opioid peptide family.

The discovery of the ORL1 receptor presents an opportunity in drug discovery for novel compounds which can be administered for pain management or other syndromes modulated by this receptor.

All documents cited herein, including the foregoing, are incorporated by reference in their entireties for all purposes.

OBJECTS AND SUMMARY OF THE INVENTION

It is accordingly an object of certain embodiments of the present invention to provide new compounds which exhibit affinity for the ORL1 receptor.

It is an object of certain embodiments of the present invention to provide new compounds which exhibit affinity for the ORL1 receptor and one or more of the μ , δ or κ receptors.

It is an object of certain embodiments of the present invention to provide new compounds for treating a patient suffering from chronic or acute pain by administering a compound having affinity for the ORL1 receptor.

It is an object of certain embodiments of the present invention to provide new compounds which have agonist activity at the μ , δ and κ receptors which is greater than compounds currently available e.g. morphine.

It is an object of certain embodiments of the present invention to provide methods of treating chronic and acute pain by administering compounds which have agonist activity at the μ , δ and κ receptors which is greater than compounds currently available.


It is an object of certain embodiments of the present invention to provide methods of treating chronic and acute pain by administering non-opioid compounds which have agonist activity at the μ , δ and κ receptors and which produce less side effects than compounds currently available.

It is an object of certain embodiments of the present invention to provide compounds useful as analgesics, anti-inflammatories, diuretics, anesthetics and neuroprotective agents, anti-hypertensives, anti-anxiolytics; agents for appetite control; hearing regulators; anti-tussives, anti-asthmatics, modulators of locomotor activity, modulators of learning and memory, regulators of neurotransmitter and hormone release, kidney function modulators, anti-depressants, agents to treat memory loss due to Alzheimer's disease or other dementias, anti-epileptics, anti-convulsants, agents to treat withdrawal from alcohol and drugs of addiction, agents to control water balance, agents to control sodium excretion and agents to control arterial blood pressure disorders and methods for administering said compounds.

The compounds of the present invention are useful for modulating a pharmacodynamic response from one or more opioid receptors (ORL-1, μ , δ and κ) centrally and/or peripherally. The response can be attributed to the compound stimulating

(agonist) or inhibiting (antagonist) the one or more receptors. Certain compounds can stimulate one receptor (e.g., a μ agonist) and inhibit a different receptor (e.g., an ORL-1 antagonist).

Other objects and advantages of the present invention will become apparent from the following detailed description thereof. The present invention in certain embodiments comprises compounds having the general formula (I):



(I)

wherein

D is a 5-8 membered cycloalkyl, 5-8 membered heterocyclic or a 6 membered aromatic or heteroaromatic group;


n is an integer from 0 to 3;

A, B and Q are independently hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{1-10} alkoxy, C_{3-12} cycloalkoxy, $-CH_2OH$, $-NHSO_2$, hydroxy C_{1-10} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, di C_{1-4} alkylaminocarbonyl-, acylamino-, acylaminoalkyl-, amide, sulfonylamino C_{1-10} alkyl-, or A-B can together form a C_{2-6} bridge, or B-Q can together form a C_{3-7} bridge, or A-Q can together form a C_{1-5} bridge;

Z is selected from the group consisting of a bond, straight or branched C_{1-6} alkylene, $-NH-$, $-CH_2O-$, $-CH_2NH-$, $-CH_2N(CH_3)-$, $-NHCH_2-$, $-CH_2CONH-$, $-NHCH_2CO-$, $-CH_2CO-$, $-COCH_2-$, $-CH_2COCH_2-$, $-CH(CH_3)-$, $-CH=$, $-O-$ and $-HC=CH-$, wherein the

carbon and/or nitrogen atoms are unsubstituted or substituted with one or more lower alkyl, hydroxy, halo or alkoxy group;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, $-COOV_1$, $-C_{1-4}COOV_1$, cyano, cyano C_{1-10} alkyl-, cyano C_{3-10} cycloalkyl-, NH_2SO_2 -, $NH_2SO_2C_{1-4}$ alkyl-, NH_2SOC_{1-4} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, di C_{1-4} alkylaminocarbonyl-, benzyl, C_{3-12} cycloalkenyl-, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a hetero-monocyclic ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ; and wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, or benzyl of R_1 is optionally substituted with 1-3 substituents selected from the group consisting of halogen, hydroxy, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, cyano, $-COOV_1$, $-C_{1-4}COOV_1$, cyano C_{1-10} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, a 5-membered heteroaromatic C_{0-4} alkyl-, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl-, C_{1-10} alkoxy-, and cyano; and wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, or spiro ring system of the formula (V) is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy or benzyloxy is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;


W_1 is hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{1-10} alkoxy, C_{3-12} cycloalkoxy, - CH_2OH , amino, C_{1-4} alkylamino-, di C_{1-4} alkylamino-, or a 5-membered heteroaromatic ring optionally substituted with 1-3 lower alkyl;

V_1 is H, C_{1-6} alkyl, C_{3-6} cycloalkyl, benzyl or phenyl;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl- and halogen, said alkyl or cycloalkyl optionally substituted with an oxo, amino, alkylamino or dialkylamino group;

and pharmaceutically acceptable salts thereof and solvates thereof.

The present invention in certain embodiments comprises compounds having the general formula (IA) as follows:




wherein

n is an integer from 0 to 3;

Z is selected from the group consisting of a bond, $-CH_2-$, $-NH-$, $-CH_2O-$, $-CH_2CH_2-$, $-CH_2NH-$, $-CH_2N(CH_3)-$, $-NHCH_2-$, $-CH_2CONH-$, $-NHCH_2CO-$, $-CH_2CO-$, $-COCH_2-$, $-CH_2COCH_2-$, $-CH(CH_3)-$, $-CH=$, and $-HC=CH-$, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with a lower alkyl, halogen, hydroxy or alkoxy group;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, benzyl, C_{3-12} cycloalkenyl, a

monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a hetero-monocyclic ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ;

wherein said monocyclic aryl is preferably phenyl;

wherein said bicyclic aryl is preferably naphthyl;

wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, or benzyl is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, cyano, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, and spiro ring system of the formula (V) are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy and benzyloxy are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

and pharmaceutically acceptable salts thereof and solvates thereof.

In certain preferred embodiments of formula (I), D is phenyl or a 6 membered heteroaromatic group containing 1-3 nitrogen atoms.

In certain preferred embodiments of formula (I) or (IA), the R_1 alkyl is methyl, ethyl, propyl, butyl, pentyl, or hexyl.

In certain preferred embodiments of formula (I) or (IA), the R_1 cycloalkyl is

cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, or norbornyl.

In other preferred embodiments of formula (I) or (IA), the R_1 bicyclic ring system is naphthyl. In other preferred embodiments of formula (I) or (IA), the R_1 bicyclic ring system is tetrahydronaphthyl, or decahydronaphthyl and the R_1 tricyclic ring system is dibenzocycloheptyl. In other preferred embodiments R_1 is phenyl or benzyl.

In other preferred embodiments of formula (I) or (IA), the R_1 bicyclic aromatic ring is a 10-membered ring, preferably quinoline or naphthyl.

In other preferred embodiments of formula (I) or (IA), the R_1 bicyclic aromatic ring is a 9-membered ring, preferably indenyl.

In certain embodiments of formula (I) or (IA), Z is a bond, methyl, or ethyl.

In certain embodiments of formula (I) or (IA), the Z group is maximally substituted as not to have any hydrogen substitution on the base Z group. For example, if the base Z group is $-\text{CH}_2-$, substitution with two methyl groups would remove hydrogens from the $-\text{CH}_2-$ base Z group.

In other preferred embodiments of formula (I) or (IA), n is 0.


In certain embodiments of formula (I) or (IA), X_1 and X_2 are both O.

In certain embodiments of formula (I), ZR_1 is cyclohexylethyl-, cyclohexylmethyl-, cyclopentylmethyl-, dimethylcyclohexylmethyl-, phenylethyl-, pyrrolyltrifluoroethyl-, thienyltrifluoroethyl-, pyridylethyl-, cyclopentyl-, cyclohexyl-, methoxycyclohexyl-, tetrahydropyranyl-, propylpiperidinyl-, indolylmethyl-, pyrazolypentyl-, thiazolyethyl-, phenyltrifluoroethyl-, hydroxyhexyl-, methoxyhexyl-, isopropoxybutyl-, hexyl-, or oxocanylpropyl-.

In certain embodiments of formula (I), ZR_1 is $-\text{CH}_2\text{COOV}_1$, tetrazolylmethyl-, cyanomethyl-, $\text{NH}_2\text{SO}_2\text{methyl-}$, $\text{NH}_2\text{SOMethyl-}$, aminocarbonylmethyl-, $\text{C}_{1-4}\text{alkylaminocarbonylmethyl-}$, or $\text{diC}_{1-4}\text{alkylaminocarbonylmethyl-}$.

In certain embodiments of formula (I), ZR_1 is 3,3 diphenylpropyl optionally substituted at the 3 carbon of the propyl with $-\text{COOV}_1$, tetrazolyl $\text{C}_{0-4}\text{alkyl-}$, cyano-, aminocarbonyl-, $\text{C}_{1-4}\text{alkylaminocarbonyl-}$, or $\text{diC}_{1-4}\text{alkylaminocarbonyl-}$.

The present invention in certain embodiments comprises compounds having the general formula (II):



(II)

wherein

the dotted line represents an optional double bond;

R is hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkylC₁₋₄alkyl-, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy-, C₁₋₁₀ alkenyl, C₁₋₁₀ alkylidene, oxo, C₁₋₁₀ alkyl substituted with 1-3 halogen, C₃₋₁₂ cycloalkyl substituted with 1-3 halogen, C₃₋₁₂ cycloalkylC₁₋₄alkyl- substituted with 1-3 halogen, C₁₋₁₀ alkoxy substituted with 1-3 halogen, C₃₋₁₂ cycloalkoxy- substituted with 1-3 halogen, -COOV₁, -C₁₋₄COOV₁, -CH₂OH, -SO₂N(V₁)₂, hydroxyC₁₋₁₀alkyl-, hydroxyC₃₋₁₀cycloalkyl-, cyanoC₁₋₁₀alkyl-, cyanoC₃₋₁₀cycloalkyl-, -CON(V₁)₂, NH₂SO₂C₁₋₄alkyl-, NH₂SOC₁₋₄alkyl-, sulfonylaminoC₁₋₁₀alkyl-, diaminoalkyl-, -sulfonylC₁₋₄alkyl, a 6-membered heterocyclic ring, a 6-membered heteroaromatic ring, a 6-membered heterocyclicC₁₋₄alkyl-, a 6-membered heteroaromaticC₁₋₄alkyl-, a 6-membered aromatic ring, a 6-membered aromaticC₁₋₄alkyl-, a 5-membered heterocyclic ring optionally substituted with an oxo or thio, a 5-membered heteroaromatic ring, a 5-membered heterocyclicC₁₋₄alkyl- optionally substituted with an oxo or thio, a 5-membered heteroaromaticC₁₋₄alkyl-, -C₁₋₅(=O)W₁, -C₁₋₅(=NH)W₁, -C₁₋₅NHC(=O)W₁, -C₁₋₅NHS(=O)₂W₁, -C₁₋₅NHS(=O)W₁, wherein W₁ is hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy, -CH₂OH, amino, C₁₋₄alkylamino-, diC₁₋₄alkylamino-, or a 5-

membered heteroaromatic ring optionally substituted with 1-3 lower alkyl;

wherein each V_1 is independently selected from H, C_{1-6} alkyl, C_{3-6} cycloalkyl, benzyl and phenyl;

n is an integer from 0 to 3;


D is a 5-8 membered cycloalkyl, 5-8 membered heterocyclic or a 6 membered aromatic or heteroaromatic group;

n is an integer from 0 to 3;

A, B and Q are independently hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{1-10} alkoxy, C_{3-12} cycloalkoxy, C_{1-10} alkenyl, C_{1-10} alkylidene, oxo, $-CH_2OH$, $-NHSO_2$, hydroxy C_{1-10} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, di C_{1-4} alkylaminocarbonyl-, acylamino-, acylaminoalkyl-, amide, sulfonylamino C_{1-10} alkyl-, or A-B can together form a C_{2-6} bridge, or B-Q can together form a C_{3-7} bridge, or A-Q can together form a C_{1-5} bridge;

Z is selected from the group consisting of a bond, straight or branched C_{1-6} alkylene, $-NH-$, $-CH_2O-$, $-CH_2NH-$, $-CH_2N(CH_3)-$, $-NHCH_2-$, $-CH_2CONH-$, $-NHCH_2CO-$, $-CH_2CO-$, $-COCH_2-$, $-CH_2COCH_2-$, $-CH(CH_3)-$, $-CH=$, $-O-$ and $-HC=CH-$, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with one or more lower alkyl, hydroxy, halo or alkoxy group;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, $-COOV_1$, $-C_{1-4}COOV_1$, cyano, cyano C_{1-10} alkyl-, cyano C_{3-10} cycloalkyl-, NH_2SO_2 -, $NH_2SO_2C_{1-4}$ alkyl-, NH_2SOC_{1-4} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, di C_{1-4} alkylaminocarbonyl-, benzyl, C_{3-12} cycloalkenyl-, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a hetero-monocyclic ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):



(V)


wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ; and wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino-, C_3

$_{12}$ cycloalkylamino-, or benzyl of R_1 is optionally substituted with 1-3 substituents selected from the group consisting of halogen, hydroxy, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, cyano, $-COOV_1$, $-C_{1-4}COOV_1$, cyano C_{1-10} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, a 5-membered heteroaromatic C_{0-4} alkyl-, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl-, C_{1-10} alkoxy-, and cyano; and wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, or spiro ring system of the formula (V) is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy or benzyloxy is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl- and halogen, said alkyl or cycloalkyl optionally substituted with an oxo, amino, alkylamino or dialkylamino group;

and pharmaceutically acceptable salts thereof and solvates thereof.

The present invention in certain embodiments comprises compounds having the formula (IIA):



(IIA)


wherein

the dotted line represents an optional double bond;

Z is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{NH}-$, $-\text{CH}_2\text{O}-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{NH}-$, $-\text{CH}_2\text{N}(\text{CH}_3)-$, $-\text{NHCH}_2-$, $-\text{CH}_2\text{CONH}-$, $-\text{NHCH}_2\text{CO}-$, $-\text{CH}_2\text{CO}-$, $-\text{COCH}_2-$, $-\text{CH}_2\text{COCH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{CH}=\text{}$, and $-\text{HC}=\text{CH}-$, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with a lower alkyl, halogen, hydroxy or alkoxy group;

R and Q are the same or different and are each selected from the group consisting of hydrogen, halogen, C_{1-10} alkyl, C_{1-10} alkenyl, C_{1-10} alkylidene, C_{3-12} cycloalkyl, C_{1-10} alkoxy, and oxo;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, benzyl, C_{3-12} cycloalkenyl, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a heteromonocyclic ring, a bicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ;

wherein said monocyclic aryl is preferably phenyl;

wherein said bicyclic aryl is preferably naphthyl;

wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, or benzyl is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, cyano, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, heteromonocyclic ring, heterobicyclic ring system, and spiro ring system of the formula (V) are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy and benzyloxy

are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, and cyano;

R₂ is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

and pharmaceutically acceptable salts thereof.

In certain preferred embodiments Q of formula (II) or (IIA), is hydrogen or methyl.

In certain preferred embodiments, R of formula (II) or (IIA), is hydrogen, methyl, ethyl, or ethylidene.

In certain preferred embodiments of formula (II), D is phenyl or a 6 membered heteroaromatic group containing 1-3 nitrogen atoms.

In certain preferred embodiments of formula (II) or (IIA), the R₁ alkyl is methyl, ethyl, propyl, butyl, pentyl, or hexyl.

In certain preferred embodiments of formula (II) or (IIA), the R₁ cycloalkyl is cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, or norbornyl.

In other preferred embodiments of formula (II) or (IIA), the R₁ bicyclic ring system is naphthyl. In other preferred embodiments of formula (II) or (IIA), the R₁ bicyclic ring system is tetrahydronaphthyl, or decahydronaphthyl and the R₁ tricyclic ring system is dibenzocycloheptyl. In other preferred embodiments R₁ is phenyl or benzyl.

In other preferred embodiments of formula (II) or (IIA), the R₁ bicyclic aromatic ring is a 10-membered ring, preferably quinoline or naphthyl.

In other preferred embodiments of formula (II) or (IIA), the R₁ bicyclic aromatic ring is a 9-membered ring, preferably indenyl.

In certain embodiments of formula (II) or (IIA), Z is a bond, methyl, or ethyl.

In certain embodiments of formula (II) or (IIA), the Z group is maximally substituted as not to have any hydrogen substitution on the base Z group. For example, if the base Z group is -CH₂-, substitution with two methyl groups would remove hydrogens from the -CH₂- base Z group.

In other preferred embodiments of formula (II) or (IIA), n is 0.

In certain embodiments of formula (II) or (IIA), X₁ and X₂ are both O.

In other preferred embodiments, the dotted line is a double bond.

In certain embodiments of formula (II), R is -CH₂C(=O)NH₂, -C(NH)NH₂, pyridylmethyl, cyclopentyl, cyclohexyl, furanylmethyl, -C(=O)CH₃, -CH₂CH₂NHC(=O)CH₃, -SO₂CH₃, CH₂CH₂NHSO₂CH₃, furanylethyl, methylpyrrolylcarbonyl-, diazolecarbonyl-, azolemethyl-, trifluoroethyl-, hydroxyethyl-, cyanomethyl-, oxo-oxazolemethyl-, or diazolemethyl-.


In certain embodiments of formula (II), ZR₁ is cyclohexylethyl-, cyclohexylmethyl-, cyclopentylmethyl-, dimethylcyclohexylmethyl-, phenylethyl-, pyrrolyltrifluoroethyl-,

thienyltrifluoroethyl-, pyridylethyl-, cyclopentyl-, cyclohexyl-, methoxycyclohexyl-, tetrahydropyranyl-, propylpiperidiny-, indolylmethyl-, pyrazolypentyl-, thiazolyethyl-, phenyltrifluoroethyl-, hydroxyhexyl-, methoxyhexyl-, isopropoxybutyl-, hexyl-, or oxocanylpropyl-.

In certain embodiments of formula (II), at least one of ZR_1 or R is $-CH_2COOV_1$, tetrazolylmethyl-, cyanomethyl-, NH_2SO_2 methyl-, NH_2SO methyl-, aminocarbonylmethyl-, C_{1-4} alkylaminocarbonylmethyl-, or di C_{1-4} alkylaminocarbonylmethyl-.

In certain embodiments of formula (II), ZR_1 is 3,3 diphenylpropyl optionally substituted at the 3 carbon of the propyl with $-COOV_1$, tetrazolyl C_{0-4} alkyl-, cyano-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, or di C_{1-4} alkylaminocarbonyl-.

The present invention in certain embodiments comprises compounds having the general formula (III):



(III)

wherein R is hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{3-12} cycloalkyl C_{1-4} alkyl-, C_{1-10} alkoxy, C_{3-12} cycloalkoxy-, C_{1-10} alkyl substituted with 1-3 halogen, C_{3-12} cycloalkyl substituted with 1-3 halogen, C_{3-12} cycloalkyl C_{1-4} alkyl- substituted with 1-3 halogen, C_{1-10} alkoxy substituted with 1-3 halogen, C_{3-12} cycloalkoxy- substituted with 1-3 halogen, $-COOV_1$, $-C_{1-4}COOV_1$, $-CH_2OH$, $-SO_2N(V_1)_2$, hydroxy C_{1-10} alkyl-, hydroxy C_{3-10} cycloalkyl-, cyano C_{1-10} alkyl-, cyano C_{3-10} cycloalkyl-, $-CON(V_1)_2$, $NH_2SO_2C_{1-4}$ alkyl-, NH_2SOC_{1-4} alkyl-, sulfonylamino C_{1-10} alkyl-, diaminoalkyl-, -sulfonyl C_{1-4} alkyl, a 6-membered heterocyclic ring, a 6-membered heteroaromatic ring, a 6-membered heterocyclic C_{1-4} alkyl-, a 6-membered heteroaromatic C_{1-4} alkyl-, a 6-membered aromatic ring, a 6-membered aromatic C_{1-4} alkyl-,


a 5-membered heterocyclic ring optionally substituted with an oxo or thio, a 5-membered heteroaromatic ring, a 5-membered heterocyclic C_{1-4} alkyl- optionally substituted with an oxo or thio, a 5-membered heteroaromatic C_{1-4} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}(=NH)W_1$, $-C_{1-5}NHC(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, wherein W_1 is hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{1-10} alkoxy, C_{3-12} cycloalkoxy, $-CH_2OH$, amino, C_{1-4} alkylamino-, di C_{1-4} alkylamino-, or a 5-membered heteroaromatic ring optionally substituted with 1-3 lower alkyl;

wherein each V_1 is independently selected from H, C_{1-6} alkyl, C_{3-6} cycloalkyl, benzyl and phenyl;

n is an integer from 0 to 3;

D is a 5-8 membered cycloalkyl, 5-8 membered heterocyclic or a 6 membered aromatic or heteroaromatic group;

Z is selected from the group consisting of a bond, straight or branched C_{1-6} alkylene, $-NH-$, $-CH_2O-$, $-CH_2NH-$, $-CH_2N(CH_3)-$, $-NHCH_2-$, $-CH_2CONH-$, $-NHCH_2CO-$, $-CH_2CO-$, $-COCH_2-$, $-CH_2COCH_2-$, $-CH(CH_3)-$, $-CH=$, $-O-$ and $-HC=CH-$, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with one or more lower alkyl, hydroxy, halo or alkoxy group; or Z is a cycloalkylamino system of the formula (VI):




(VI)

wherein A, B and Q are independently hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{1-10} alkoxy, C_{3-12} cycloalkoxy, $-CH_2OH$, $-NHSO_2$, hydroxy C_{1-10} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, di C_{1-4} alkylaminocarbonyl-, acylamino-, acylaminoalkyl-, amide, sulfonylamino C_{1-10} alkyl-, or A-B can together form a C_{2-6} bridge, or B-Q can together form a C_{3-7} bridge, or A-Q can together form a C_{1-5} bridge;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, $-COOV_1$, $-C_{1-4}COOV_1$, cyano, cyano C_{1-10} alkyl-, cyano C_{3-10} cycloalkyl-, NH_2SO_2 , $NH_2SO_2C_{1-4}$ alkyl-, NH_2SOC_{1-4} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, di C_{1-4} alkylaminocarbonyl-, benzyl, C_{3-12}

cycloalkenyl-, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a hetero-monocyclic ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):




(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ; and wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, or benzyl of R_1 is optionally substituted with 1-3 substituents selected from the group consisting of halogen, hydroxy, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, cyano, $-COOV_1$, $-C_{1-4}COOV_1$, cyano C_{1-10} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, a 5-membered heteroaromatic C_{0-4} alkyl-, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl-, C_{1-10} alkoxy-, and cyano; and wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, or spiro ring system of the formula (V) is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy or benzyloxy is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl- and halogen, said alkyl or cycloalkyl optionally substituted with an oxo, amino, alkylamino or dialkylamino group;

and pharmaceutically acceptable salts thereof and solvates thereof.

The present invention in certain embodiments comprises compounds having the formula (IIIa):




(III A)

wherein

n is an integer from 0 to 3;

Z is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{NH}-$, $-\text{CH}_2\text{O}-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{NH}-$, $-\text{CH}_2\text{N}(\text{CH}_3)-$, $-\text{NHCH}_2-$, $-\text{CH}_2\text{CONH}-$, $-\text{NHCH}_2\text{CO}-$, $-\text{CH}_2\text{CO}-$, $-\text{COCH}_2-$, $-\text{CH}_2\text{COCH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{CH}=\text{}$, $-\text{HC}=\text{CH}-$, and a cycloalkylamino system of the formula (VI):




(VI)

wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with a lower alkyl, halogen, hydroxy, phenyl, benzyl, or alkoxy group;

R is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{1-10} alkoxy, and C_{3-12} cycloalkyl;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, benzyl, C_{3-12} cycloalkenyl, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a heteromonocyclic ring, a heterobicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ;

wherein said monocyclic aryl is preferably phenyl;

wherein said bicyclic aryl is preferably naphthyl;

wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, or benzyl is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, cyano, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, heteromonocyclic ring, heterobicyclic ring system, and spiro ring system of the formula (V) are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy and benzyloxy are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

and pharmaceutically acceptable salts thereof.

In certain preferred embodiments of formula (III), D is phenyl or a 6 membered heteroaromatic group containing 1-3 nitrogen atoms.

In certain preferred embodiments of formula (III) or (IIIA), the R_1 alkyl is methyl, ethyl, propyl, butyl, pentyl, or hexyl.

In certain preferred embodiments of formula (III) or (IIIA), the R_1 cycloalkyl is cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, or norbornyl.

In other preferred embodiments of formula (III) or (IIIA), the R_1 bicyclic ring

system is naphthyl. In other preferred embodiments of formula (III) or (IIIA), the R_1 bicyclic ring system is tetrahydronaphthyl, or decahydronaphthyl and the R_1 tricyclic ring system is dibenzocycloheptyl. In other preferred embodiments R_1 is phenyl or benzyl.


In other preferred embodiments of formula (III) or (IIIA), the R_1 bicyclic aromatic ring is a 10-membered ring, preferably quinoline or naphthyl.

In other preferred embodiments of formula (III) or (IIIA), the R_1 bicyclic aromatic ring is a 9-membered ring, preferably indenyl.

In certain embodiments of formula (III) or (IIIA), Z is a bond, methyl, or ethyl.

In certain embodiments of formula (III) or (IIIA), the Z group is maximally substituted as not to have any hydrogen substitution on the base Z group. For example, if the base Z group is $-\text{CH}_2-$, substitution with two methyl groups would remove hydrogens from the $-\text{CH}_2-$ base Z group.

In certain embodiments of formula (III) or (IIIA), Z is a cycloalkylamino system of the formula (VI):



(VI)

wherein the nitrogen atom is optionally substituted with a C_{1-3} alkyl, phenyl, or benzyl.

In other preferred embodiments of formula (III) or (IIIA), n is 0.

In certain embodiments of formula (III) or (IIIA), X_1 and X_2 are both O.

In certain embodiments of formula (III), R is $-\text{CH}_2\text{C}(=\text{O})\text{NH}_2$, $-\text{C}(\text{NH})\text{NH}_2$, pyridylmethyl, cyclopentyl, cyclohexyl, furanylmethyl, $-\text{C}(=\text{O})\text{CH}_3$, $-\text{CH}_2\text{CH}_2\text{NHC}(=\text{O})\text{CH}_3$, $-\text{SO}_2\text{CH}_3$, $\text{CH}_2\text{CH}_2\text{NHSO}_2\text{CH}_3$, furanylcarbonyl-, methylpyrrolylcarbonyl-, diazolecarbonyl-, azolemethyl-, trifluoroethyl-, hydroxyethyl-, cyanomethyl-, oxo-oxazolemethyl-, or diazolemethyl-.


In certain embodiments of formula (III), ZR_1 is cyclohexylethyl-, cyclohexylmethyl-, cyclopentylmethyl-, dimethylcyclohexylmethyl-, phenylethyl-, pyrrolyltrifluoroethyl-, thienyltrifluoroethyl-, pyridylethyl-, cyclopentyl-, cyclohexyl-, methoxycyclohexyl-, tetrahydropyranyl-, propylpiperidinyl-, indolylmethyl-, pyrazoylpentyl-, thiazolyethyl-,

phenyltrifluoroethyl-, hydroxyhexyl-, methoxyhexyl-, isopropoxybutyl-, hexyl-, or oxocanylpropyl-.

In certain embodiments of formula (III), at least one of ZR_1 or R is $-CH_2COOV_1$, tetrazolylmethyl-, cyanomethyl-, NH_2SO_2 methyl-, NH_2SO methyl-, aminocarbonylmethyl-, C_{1-4} alkylaminocarbonylmethyl-, or di C_{1-4} alkylaminocarbonylmethyl-.

In certain embodiments of formula (III), ZR_1 is 3,3 diphenylpropyl optionally substituted at the 3 carbon of the propyl with $-COOV_1$, tetrazolyl C_{0-4} alkyl-, cyano-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, or di C_{1-4} alkylaminocarbonyl-.

The present invention in certain embodiments comprises compounds having the general formula (IV):



(IV)

wherein R is hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{3-12} cycloalkyl C_{1-4} alkyl-, C_{1-10} alkoxy, C_{3-12} cycloalkoxy-, C_{1-10} alkyl substituted with 1-3 halogen, C_{3-12} cycloalkyl substituted with 1-3 halogen, C_{3-12} cycloalkyl C_{1-4} alkyl- substituted with 1-3 halogen, C_{1-10} alkoxy substituted with 1-3 halogen, C_{3-12} cycloalkoxy- substituted with 1-3 halogen, $-COOV_1$, $-C_{1-4}COOV_1$, $-CH_2OH$, $-SO_2N(V_1)_2$, hydroxy C_{1-10} alkyl-, hydroxy C_{3-10} cycloalkyl-, cyano C_{1-10} alkyl-, cyano C_{3-10} cycloalkyl-, $-CON(V_1)_2$, $NH_2SO_2C_{1-4}$ alkyl-, NH_2SOC_{1-4} alkyl-, sulfonylamino C_{1-10} alkyl-, diaminoalkyl-, -sulfonyl C_{1-4} alkyl, a 6-membered heterocyclic ring, a 6-membered heteroaromatic ring, a 6-membered heterocyclic C_{1-4} alkyl-, a 6-membered heteroaromatic C_{1-4} alkyl-, a 6-membered aromatic ring, a 6-membered aromatic C_{1-4} alkyl-,

a 5-membered heterocyclic ring optionally substituted with an oxo or thio, a 5-membered heteroaromatic ring, a 5-membered heterocyclicC₁₋₄alkyl- optionally substituted with an oxo or thio, a 5-membered heteroaromaticC₁₋₄alkyl-, -C₁₋₅(=O)W₁, -C₁₋₅(=NH)W₁, -C₁₋₅NHC(=O)W₁, -C₁₋₅NHS(=O)₂W₁, -C₁₋₅NHS(=O)W₁, wherein W₁ is hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy, -CH₂OH, amino, C₁₋₄alkylamino-, diC₁₋₄alkylamino-, or a 5-membered heteroaromatic ring optionally substituted with 1-3 lower alkyl;

wherein each V₁ is independently selected from H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, benzyl and phenyl;


D is a 5-8 membered cycloalkyl, 5-8 membered heterocyclic or a 6 membered aromatic or heteroaromatic group;

n is an integer from 0 to 3;

A, B and Q are independently hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy, -CH₂OH, -NHSO₂, hydroxyC₁₋₁₀alkyl-, aminocarbonyl-, C₁₋₄alkylaminocarbonyl-, diC₁₋₄alkylaminocarbonyl-, acylamino-, acylaminoalkyl-, amide, sulfonylaminoC₁₋₁₀alkyl-, or A-B can together form a C₂₋₆ bridge, or B-Q can together form a C₃₋₇ bridge, or A-Q can together form a C₁₋₅ bridge;

Z is selected from the group consisting of a bond, straight or branched C₁₋₆ alkylene, -NH-, -CH₂O-, -CH₂NH-, -CH₂N(CH₃)-, -NHCH₂-, -CH₂CONH-, -NHCH₂CO-, -CH₂CO-, -COCH₂-, -CH₂COCH₂-, -CH(CH₃)-, -CH=, -O- and -HC=CH-, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with one or more lower alkyl, hydroxy, halo or alkoxy group;

R₁ is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂cycloalkyl, C₂₋₁₀alkenyl, amino, C₁₋₁₀alkylamino-, C₃₋₁₂cycloalkylamino-, -COOV₁, -C₁₋₄COOV₁, cyano, cyanoC₁₋₁₀alkyl-, cyanoC₃₋₁₀cycloalkyl-, NH₂SO₂-, NH₂SO₂C₁₋₄alkyl-, NH₂SOC₁₋₄alkyl-, aminocarbonyl-, C₁₋₄alkylaminocarbonyl-, diC₁₋₄alkylaminocarbonyl-, benzyl, C₃₋₁₂cycloalkenyl-, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a heteromonocyclic ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):




(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ; and wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, or benzyl of R_1 is optionally substituted with 1-3 substituents selected from the group consisting of halogen, hydroxy, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, cyano, $-COOV_1$, $-C_{1-4}COOV_1$, cyano C_{1-10} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, a 5-membered heteroaromatic C_{0-4} alkyl-, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl-, C_{1-10} alkoxy-, and cyano; and wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, or spiro ring system of the formula (V) is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy or benzyloxy is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl- and halogen, said alkyl or cycloalkyl optionally substituted with an oxo, amino, alkylamino or dialkylamino group;

and pharmaceutically acceptable salts thereof and solvates thereof.

The present invention in certain embodiments comprises compounds having the formula (IVA):




wherein

n is an integer from 0 to 3;

Z is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{NH}-$, $-\text{CH}_2\text{O}-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{NH}-$, $-\text{CH}_2\text{N}(\text{CH}_3)-$, $-\text{NHCH}_2-$, $-\text{CH}_2\text{CONH}-$, $-\text{NHCH}_2\text{CO}-$, $-\text{CH}_2\text{CO}-$, $-\text{COCH}_2-$, $-\text{CH}_2\text{COCH}_2-$, $-\text{CH}(\text{CH}_3)-$, $-\text{CH}=\text{}$, and $-\text{HC}=\text{CH}-$, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with a lower alkyl, halogen, hydroxy or alkoxy group;

R is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{1-10} alkoxy, and C_{3-12} cycloalkyl;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, benzyl, C_{3-12} cycloalkenyl, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a heteromonocyclic ring, a heterobicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ;

wherein said monocyclic aryl is preferably phenyl;

wherein said bicyclic aryl is preferably naphthyl;

wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, or benzyl is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, cyano, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, heteromonocyclic ring, heterobicyclic ring system, and spiro ring system of the formula (V) are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy and benzyloxy

are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, and cyano;

R₂ is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

and pharmaceutically acceptable salts thereof.

In certain preferred embodiments of formula (IV), D is phenyl or a 6 membered heteroaromatic group containing 1-3 nitrogen atoms.

In certain preferred embodiments of formula (IV) or (IVA), the R₁ alkyl is methyl, ethyl, propyl, butyl, pentyl, or hexyl.

In certain preferred embodiments of formula (IV) or (IVA), the R₁ cycloalkyl is cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, or norbornyl.

In other preferred embodiments of formula (IV) or (IVA), the R₁ bicyclic ring system is naphthyl. In other preferred embodiments of formula (IV) or (IVA), the R₁ bicyclic ring system is tetrahydronaphthyl, or decahydronaphthyl and the R₁ tricyclic ring system is dibenzocycloheptyl. In other preferred embodiments R₁ is phenyl or benzyl.

In other preferred embodiments of formula (IV) or (IVA), the R₁ bicyclic aromatic ring is a 10-membered ring, preferably quinoline or naphthyl.

In other preferred embodiments of formula (IV) or (IVA), the R₁ bicyclic aromatic ring is a 9-membered ring, preferably indenyl.

In certain embodiments of formula (IV) or (IVA), Z is a bond, methyl, or ethyl.

In certain embodiments of formula (IV) or (IVA), the Z group is maximally substituted as not to have any hydrogen substitution on the base Z group. For example, if the base Z group is -CH₂-, substitution with two methyl groups would remove hydrogens from the -CH₂- base Z group.

In other preferred embodiments of formula (IV) or (IVA), n is 0.

In certain embodiments of formula (IV) or (IVA), X₁ and X₂ are both O.

In certain embodiments of formula (IV), R is -CH₂C(=O)NH₂, -C(NH)NH₂, pyridylmethyl, cyclopentyl, cyclohexyl, furanylmethyl, -C(=O)CH₃, -CH₂CH₂NHC(=O)CH₃, -SO₂CH₃, CH₂CH₂NHSO₂CH₃, furanylethyl, methylpyrrolylcarbonyl-, diazolecarbonyl-, azolemethyl-, trifluoroethyl-, hydroxyethyl-, cyanomethyl-, oxo-oxazolemethyl-, or diazolemethyl-.


In certain embodiments of formula (IV), ZR₁ is cyclohexylethyl-, cyclohexylmethyl-, cyclopentylmethyl-, dimethylcyclohexylmethyl-, phenylethyl-, pyrrolyltrifluoroethyl-, thienyltrifluoroethyl-, pyridylethyl-, cyclopentyl-, cyclohexyl-, methoxycyclohexyl-, tetrahydropyranyl-, propylpiperidinyl-, indolylmethyl-, pyrazoylpentyl-, thiazolyethyl-, phenyltrifluoroethyl-, hydroxyhexyl-, methoxyhexyl-, isopropoxybutyl-, hexyl-, or

oxocanylpropyl-.

In certain embodiments of formula (IV), at least one of ZR_1 or R is $-CH_2COOV_1$, tetrazolylmethyl-, cyanomethyl-, NH_2SO_2 methyl-, NH_2SO methyl-, aminocarbonylmethyl-, C_{1-4} alkylaminocarbonylmethyl-, or diC_{1-4} alkylaminocarbonylmethyl-.


In certain embodiments of formula (IV), ZR_1 is 3,3 diphenylpropyl optionally substituted at the 3 carbon of the propyl with $-COOV_1$, tetrazolyl C_{0-4} alkyl-, cyano-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, or diC_{1-4} alkylaminocarbonyl-.

In alternate embodiments of formulae (I), (IA), (II), (IIA), (III), (IIIA), (IV), and (IVA), ZR_1 can be the following



wherein

Y_1 is $R_3-(C_1-C_{12})$ alkyl, R_4 -aryl, R_5 -heteroaryl, $R_6-(C_3-C_{12})$ cyclo-alkyl, $R_7-(C_3-C_7)$ heterocycloalkyl, $-CO_2(C_1-C_6)$ alkyl, CN or $-C(O)NR_8R_9$; Y_2 is hydrogen or Y_1 ; Y_3 is hydrogen or (C_1-C_6) alkyl; or Y_1 , Y_2 and Y_3 , together with the carbon to which they are attached, form one of the following structures:



wherein r is 0 to 3; w and u are each 0-3, provided that the sum of w and u is 1-3; c and d are independently 1 or 2; s is 1 to 5; and ring E is a fused R_4 -phenyl or R_5 -heteroaryl ring;

R_{10} is 1 to 3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, $-OR_8$, $-(C_1-C_6)alkyl-OR_8$, $-NR_8R_9$ and $-(C_1-C_6)alkyl-NR_8R_9$;


R_{11} is 1 to 3 substituents independently selected from the group consisting of R_{10} , $-CF_3$, $-OCF_3$, NO_2 and halo, or R_{11} substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;

R_8 and R_9 are independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_3-C_{12}) cycloalkyl, aryl and aryl (C_1-C_6) alkyl;

R_3 is 1 to 3 substituents independently selected from the group consisting of H, R_4 -aryl, R_6 -(C_3-C_{12})cycloalkyl, R_5 -heteroaryl, R_7 -(C_3-C_7)heterocycloalkyl, $-NR_8R_9$, $-OR_{12}$ and $-S(O)_{0-2}R_{12}$;

R_6 is 1 to 3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, R_4 -aryl, $-NR_8R_9$, $-OR_{12}$ and $-SR_{12}$;

R_4 is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C_1-C_6) alkyl, R_{13} -aryl, (C_3-C_{12}) cycloalkyl, $-CN$, $-CF_3$, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-OCF_3$, $-NR_8R_9$, $-(C_1-C_6)$ alkyl- NR_8R_9 , $-NHSO_2R_8$, $-SO_2N(R_{14})_2$, $-SO_2R_8$, $-SOR_8$, $-SR_8$, $-NO_2$, $-CONR_8R_9$, $-NR_9COR_8$, $-COR_8$, $-COCF_3$, $-OCOR_8$, $-OCO_2R_8$, $-COOR_8$, $-(C_1-C_6)$ alkyl- $NHCOOC(CH_3)_3$, $-(C_1-C_6)$ alkyl- $NHCOCF_3$, $-(C_1-C_6)$ alkyl- $NHSO_2$ - (C_1-C_6) alkyl, $-(C_1-C_6)$ alkyl- $NHCONH$ - (C_1-C_6) alkyl and



wherein f is 0 to 6; or R_4 substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;

R_5 is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C_1-C_6) alkyl, R_{13} -aryl, (C_3-C_{12}) cycloalkyl, $-CN$, $-CF_3$, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-OCF_3$, $-NR_8R_9$, $-(C_1-C_6)$ alkyl- NR_8R_9 , $-NHSO_2R_8$, $-SO_2N(R_{14})_2$, $-NO_2$, $-CONR_8R_9$, $-NR_9COR_8$, $-COR_8$, $-OCOR_8$, $-OCO_2R_8$ and $-COOR_8$;

R_7 is H, (C_1-C_6) alkyl, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-NR_8R_9$ or $-(C_1-C_6)$ alkyl- NR_8R_9 ;

R_{12} is H, (C_1-C_6) alkyl, R_4 -aryl, $-(C_1-C_6)$ alkyl- OR_8 , $-(C_1-C_6)$ alkyl- NR_8R_9 , $-(C_1-C_6)$ alkyl- SR_8 , or aryl (C_1-C_6) alkyl;

R_{13} is 1-3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, (C_1-C_6) alkoxy and halo;

R_{14} is independently selected from the group consisting of H, (C_1-C_6) alkyl and R_{13} - $C_6H_4-CH_2$ -.

As used herein, the term "alkyl" means a linear or branched saturated aliphatic hydrocarbon group having a single radical and 1-10 carbon atoms. Examples of alkyl groups include methyl, propyl, isopropyl, butyl, n-butyl, isobutyl, sec-butyl, tert-butyl, and pentyl. A branched alkyl means that one or more alkyl groups such as methyl, ethyl or propyl, replace one or both hydrogens in a $-CH_2-$ group of a linear alkyl chain. The term "lower alkyl" means an alkyl of 1-3 carbon atoms.

The term "alkoxy" means an "alkyl" as defined above connected to an oxygen radical.

The term "cycloalkyl" means a non-aromatic mono- or multicyclic hydrocarbon ring system having a single radical and 3-12 carbon atoms. Exemplary monocyclic

cycloalkyl rings include cyclopropyl, cyclopentyl, and cyclohexyl. Exemplary multicyclic cycloalkyl rings include adamantyl and norbornyl.

The term "alkenyl" means a linear or branched aliphatic hydrocarbon group containing a carbon-carbon double bond having a single radical and 2-10 carbon atoms.

A "branched" alkenyl means that one or more alkyl groups such as methyl, ethyl or propyl replace one or both hydrogens in a $-CH_2-$ or $-CH=$ linear alkenyl chain. Exemplary alkenyl groups include ethenyl, 1- and 2- propenyl, 1-, 2- and 3- butenyl, 3-methylbut-2-enyl, 2-propenyl, heptenyl, octenyl and decenyl.

The term "cycloalkenyl" means a non-aromatic monocyclic or multicyclic hydrocarbon ring system containing a carbon-carbon double bond having a single radical and 3 to 12 carbon atoms. Exemplary monocyclic cycloalkenyl rings include cyclopropenyl, cyclopentenyl, cyclohexenyl or cycloheptenyl. An exemplary multicyclic cycloalkenyl ring is norbornenyl.

The term "aryl" means a carbocyclic aromatic ring system containing one, two or three rings which may be attached together in a pendent manner or fused, and containing a single radical. Exemplary aryl groups include phenyl, naphthyl and acenaphthyl.

The term "heterocyclic" means cyclic compounds having one or more heteroatoms (atoms other than carbon) in the ring, and having a single radical. The ring may be saturated, partially saturated or unsaturated, and the heteroatoms may be selected from the group consisting of nitrogen, sulfur and oxygen. Examples of saturated heterocyclic radicals include saturated 3 to 6- membered hetero-monocyclic groups containing 1 to 4 nitrogen atoms, such as pyrrolidinyl, imidazolidinyl, piperidino, piperazinyl; saturated 3- to 6- membered hetero-monocyclic groups containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, such as morpholinyl; saturated 3- to 6- membered hetero-monocyclic groups containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, such as thiazolidinyl. Examples of partially saturated heterocyclic radicals include dihydrothiophene, dihydropyran, and dihydrofuran. Other heterocyclic groups can be 7 to 10 carbon rings substituted with heteroatoms such as oxocanyl and thiocanyl. When the heteroatom is sulfur, the sulfur can be a sulfur dioxide such as thiocanyldioxide.

The term "heteroaryl" means unsaturated heterocyclic radicals, wherein "heterocyclic" is as previously described. Exemplary heteroaryl groups include unsaturated 3 to 6 membered hetero-monocyclic groups containing 1 to 4 nitrogen atoms, such as pyrrolyl, pyridyl, pyrimidyl, and pyrazinyl; unsaturated condensed heterocyclic groups containing 1 to 5 nitrogen atoms, such as indolyl, quinolyl and isoquinolyl;

unsaturated 3 to 6-membered hetero-monocyclic groups containing an oxygen atom, such as furyl; unsaturated 3 to 6 membered hetero-monocyclic groups containing a sulfur atom, such as thienyl; unsaturated 3 to 6 membered hetero-monocyclic groups containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, such as oxazolyl; unsaturated condensed heterocyclic groups containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, such as benzoxazolyl; unsaturated 3 to 6 membered hetero-monocyclic groups containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, such as thiazolyl; and unsaturated condensed heterocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, such as benzothiazolyl. The term "heteroaryl" also includes unsaturated heterocyclic radicals, wherein "heterocyclic" is as previously described, in which the heterocyclic group is fused with an aryl group, in which aryl is as previously described. Exemplary fused radicals include benzofuran, benzodioxole and benzothiophene.

As used herein, the term "heterocyclic C_{1-4} alkyl", "heteroaromatic C_{1-4} alkyl" and the like refer to the ring structure bonded to a C_{1-4} alkyl radical.

All of the cyclic ring structures disclosed herein can be attached at any point where such connection is possible, as recognized by one skilled in the art.

As used herein, the term "patient" includes a human or an animal such as a companion animal or livestock.

As used herein, the term "halogen" includes fluoride, bromide, chloride, iodide or alabamide.

The invention disclosed herein is meant to encompass all pharmaceutically acceptable salts thereof of the disclosed compounds. The pharmaceutically acceptable salts include, but are not limited to, metal salts such as sodium salt, potassium salt, cesium salt and the like; alkaline earth metals such as calcium salt, magnesium salt and the like; organic amine salts such as triethylamine salt, pyridine salt, picoline salt, ethanolamine salt, triethanolamine salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt and the like; inorganic acid salts such as hydrochloride, hydrobromide, sulfate, phosphate and the like; organic acid salts such as formate, acetate, trifluoroacetate, maleate, fumarate, tartrate and the like; sulfonates such as methanesulfonate, benzenesulfonate, p-toluenesulfonate, and the like; amino acid salts such as arginate, aspartate, glutamate and the like.

The invention disclosed herein is also meant to encompass all prodrugs of the disclosed compounds. Prodrugs are considered to be any covalently bonded carriers which release the active parent drug in vivo.

The invention disclosed herein is also meant to encompass the in vivo metabolic

products of the disclosed compounds. Such products may result for example from the oxidation, reduction, hydrolysis, amidation, esterification and the like of the administered compound, primarily due to enzymatic processes. Accordingly, the invention includes compounds produced by a process comprising contacting a compound of this invention with a mammal for a period of time sufficient to yield a metabolic product thereof. Such products typically are identified by preparing a radiolabelled compound of the invention, administering it parenterally in a detectable dose to an animal such as rat, mouse, guinea pig, monkey, or to man, allowing sufficient time for metabolism to occur and isolating its conversion products from the urine, blood or other biological samples.

The invention disclosed herein is also meant to encompass the disclosed compounds being isotopically-labelled by having one or more atoms replaced by an atom having a different atomic mass or mass number. Examples of isotopes that can be incorporated into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Some of the compounds disclosed herein may contain one or more asymmetric centers and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms. The present invention is also meant to encompass all such possible forms as well as their racemic and resolved forms and mixtures thereof. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended to include both E and Z geometric isomers. All tautomers are intended to be encompassed by the present invention as well.

As used herein, the term "stereoisomers" is a general term for all isomers of individual molecules that differ only in the orientation of their atoms in space. It includes enantiomers and isomers of compounds with more than one chiral center that are not mirror images of one another (diastereomers).

The term "chiral center" refers to a carbon atom to which four different groups are attached.

The term "enantiomer" or "enantiomeric" refers to a molecule that is nonsuperimposable on its mirror image and hence optically active wherein the enantiomer rotates the plane of polarized light in one direction and its mirror image rotates the plane of polarized light in the opposite direction.

The term "racemic" refers to a mixture of equal parts of enantiomers and which is optically inactive.

The term "resolution" refers to the separation or concentration or depletion of one of the two enantiomeric forms of a molecule.

The term "modulate" as used herein with respect to the ORL-1 receptor means the mediation of a pharmacodynamic response (e.g., analgesia) in a subject from (i) inhibiting or activating the receptor, or (ii) directly or indirectly affecting the normal regulation of the receptor activity. Compounds which modulate the receptor activity include agonists, antagonists, mixed agonists/antagonists and compounds which directly or indirectly affect regulation of the receptor activity.

Certain preferred compounds of formula (I) and (IA) include:

3-[1-(naphth-2-yl-methyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(naphth-1-yl-methyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(*p*-phenylbenzyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(*p*-benzyloxybenzyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(*p*-cyanobenzyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(3,3-diphenylpropyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-[4,4-Bis-(4-fluorophenyl)butyl]-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(2-phenylethyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(cyclooctylmethyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(5-methylhex-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(10,11-Dihydro-5H-dibenzo[a,d]-cyclohepten-5-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(4-propyl-cyclohexyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(norbornan-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(decahydro-2-naphthyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(3,3-dimethyl-1,5-dioxaspiro[5.5]undeca-9-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-[4-(1-methylethyl)-cyclohexyl]-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 3-[1-(cyclooctyl)-4-piperidinyl]-2H-benzoxazol-2-one; and
 pharmaceutically acceptable salts thereof and solvates thereof.

Certain preferred compounds of formula (II) and (IIA) include:

3-ethylidene-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethylidene-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethylidene-1-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethylidene-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-

one;

3-ethylidene-1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-

one;

3-ethylidene-1-[1-(*p*-benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-

one;

3-ethylidene-1-[1-(benzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethylidene-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethylidene-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethylidene-1-[1-(3,3-diphenylpropyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-

one;

3-ethylidene-1-[1-(*p*-cyanobenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-[4-(1-methylethyl)-cyclohexyl]-4-piperidinyl]-1,3-dihydro-2H-indole-

2-one;

3-ethyl-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-

2-one;

3-ethyl-1-[1-(decahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(naphth-1-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(*p*-phenylbenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(3,3-Bis(phenyl)propyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(*p*-cyanobenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(*p*-benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(1,2,3,4-tetrahydronaphth-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(benzyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(4-propyl-cyclohexyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(5-methylhex-2-yl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

1-[1-(decahydro-2-naphthyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-

one;

1-[1-(4-(1-methylethyl)-cyclohexyl)-3-(methyl)-4-piperidiny]-1,3-dihydro-2H-indole-2-one;

1-[1-(cyclooctylmethyl)-3-(methyl)-4-piperidiny]-1,3-dihydro-2H-indole-2-one;

1-[1-(3,3-Bis(phenyl)propyl)-3-(methyl)-4-piperidiny]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(3,3-Bis(phenyl)propyl)-3-(methyl)-4-piperidiny]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(4-propylcyclohexyl)-3-(methyl)-4-piperidiny]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(5-methylhex-2-yl)-3-(methyl)-4-piperidiny]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-[4-(1-methylethyl)cyclohexyl]-3-methyl-4-piperidiny]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(decahydro-2-naphthyl)-3-(methyl)-4-piperidiny]-1,3-dihydro-2H-indole-2-one; and

pharmaceutically acceptable salts thereof and solvates thereof.

Certain preferred compounds of formula (III) and (IIIA) include:

3-ethyl-1-(*p*-phenylbenzyl)-1,3-dihydro-2H-benzimidazol-2-one;

3-ethyl-1-(5-methylhex-2-yl)-1,3-dihydro-2H-benzimidazol-2-one;

3-ethyl-1-(4-propylcyclohexyl)-1,3-dihydro-2H-benzimidazol-2-one;

3-ethyl-1-(decahydro-2-naphthyl)-1,3-dihydro-2H-benzimidazol-2-one;

3-ethyl-1-(naphth-2-yl-methyl)-1,3-dihydro-2H-benzimidazol-2-one;

1-(*p*-benzyloxybenzyl)-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;

1-benzyl-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-(benzylamino)-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;

3-ethyl-1-(naphthylmethyl)-1,3-dihydro-2H-benzimidazol-2-one;

3-ethyl-1-[5-(3-fluorophenyl)-5-(4-fluorophenyl)-hexyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-[(naphth-2-yl-methyl)ethylamino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-(norbornan-2-ylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-[[4-(1-methylethyl)-cyclohexyl]amino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-(ethylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(benzylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(indan-2-yl)benzylamino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(cyclooctylmethyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(naphth-2-yl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(*p*-benzyloxybenzyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(cyclooctylmethyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(benzylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(dibenzylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(*p*-phenylbenzyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(1,2,3,4-tetrahydronaphthyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(4-propyl-cyclohexyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(5-methylhex-2-yl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(cyclooctylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(indan-2-yl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(4-phenyl-cyclohexyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(5-methylhex-2-yl)amino]-cyclohexyl]-7-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one; and
pharmaceutically acceptable salts thereof and solvates thereof.

Other preferred compounds formula (IV) and (IVA) include:

2-cyanoimino-3-ethyl-1-[1-(*p*-phenylbenzyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(*p*-benzyloxybenzyl)-4-piperidinyl] 1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(naphth-2-yl-methyl)-4-piperidinyl] 1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-[4-(2-propyl)-cyclohexyl]-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(decahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(10,11-dihydro-5H-dibenzo[a,d]-cyclohepten-5-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(3,3-Bis(phenyl)propyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(1,2,3,4-tetrahydronaphthyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole; and

pharmaceutically acceptable salts thereof and solvates thereof.

Other preferred compounds of formula (IV) include

2-cyanoimino-3-(2-hydroxy)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-methoxycarbonylmethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-cyanomethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-

benzimidazole;

2-cyanoimino-3-butyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-(2-methanesulfonamido)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-acetomido-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-carboxymethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-(2-dimethylamino)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-1-[1-(cyclooctyl)-3-hydroxymethyl-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-7-azabenzimidazole;

2-cyanoimino-1-[1-(cyclooctyl)-2,6-ethano-4-one-4-piperidinyl]-1,3-dihydro-2H-benzimidazole; and

pharmaceutically acceptable salts thereof and solvates thereof.

The present invention also provides use of any of the disclosed compounds in the preparation of a medicament for treating pain and other disease states modulated by an opioid receptor, e.g., the ORL-1 receptor.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of the present invention can be administered to anyone requiring modulation of the opioid and ORL1 receptors. Administration may be orally, topically, by suppository, inhalation, or parenterally.

The present invention also encompasses all pharmaceutically acceptable salts of the foregoing compounds. One skilled in the art will recognize that acid addition salts of the presently claimed compounds may be prepared by reaction of the compounds with the appropriate acid via a variety of known methods.

Various oral dosage forms can be used, including such solid forms as tablets, gelcaps, capsules, caplets, granules, lozenges and bulk powders and liquid forms such as emulsions, solution and suspensions. The compounds of the present invention can be administered alone or can be combined with various pharmaceutically acceptable carriers and excipients known to those skilled in the art, including but not limited to diluents, suspending agents, solubilizers, binders, disintegrants, preservatives, coloring agents,

lubricants and the like.

When the compounds of the present invention are incorporated into oral tablets, such tablets can be compressed, tablet triturates, enteric-coated, sugar-coated, film-coated, multiply compressed or multiply layered. Liquid oral dosage forms include aqueous and nonaqueous solutions, emulsions, suspensions, and solutions and/or suspensions reconstituted from non-effervescent granules, containing suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, coloring agents, and flavoring agents. When the compounds of the present invention are to be injected parenterally, they may be, e.g., in the form of an isotonic sterile solution. Alternatively, when the compounds of the present invention are to be inhaled, they may be formulated into a dry aerosol or may be formulated into an aqueous or partially aqueous solution.

In addition, when the compounds of the present invention are incorporated into oral dosage forms, it is contemplated that such dosage forms may provide an immediate release of the compound in the gastrointestinal tract, or alternatively may provide a controlled and/or sustained release through the gastrointestinal tract. A wide variety of controlled and/or sustained release formulations are well known to those skilled in the art, and are contemplated for use in connection with the formulations of the present invention. The controlled and/or sustained release may be provided by, e.g., a coating on the oral dosage form or by incorporating the compound(s) of the invention into a controlled and/or sustained release matrix.

Specific examples of pharmaceutically acceptable carriers and excipients that may be used to formulate oral dosage forms, are described in the Handbook of Pharmaceutical Excipients, American Pharmaceutical Association (1986). Techniques and compositions for making solid oral dosage forms are described in Pharmaceutical Dosage Forms: Tablets (Lieberman, Lachman and Schwartz, editors) 2nd edition, published by Marcel Dekker, Inc. Techniques and compositions for making tablets (compressed and molded), capsules (hard and soft gelatin) and pills are also described in Remington's Pharmaceutical Sciences (Arthur Osol, editor), 1553B1593 (1980). Techniques and composition for making liquid oral dosage forms are described in Pharmaceutical Dosage Forms: Disperse Systems, (Lieberman, Rieger and Banker, editors) published by Marcel Dekker, Inc.

When the compounds of the present invention are incorporated for parenteral administration by injection (e.g., continuous infusion or bolus injection), the formulation for parenteral administration may be in the form of suspensions, solutions, emulsions in oily or aqueous vehicles, and such formulations may further comprise pharmaceutically necessary additives such as stabilizing agents, suspending agents, dispersing agents, and the like. The compounds of the invention may also be in the form of a powder for

reconstitution as an injectable formulation.

In certain embodiments, the compounds of the present invention can be used in combination with at least one other therapeutic agent. Therapeutic agents include, but are not limited to, μ -opioid agonists; non-opioid analgesics; non-steroid antiinflammatory agents; Cox-II inhibitors; antiemetics; β -adrenergic blockers; anticonvulsants; antidepressants; Ca²⁺-channel blockers; anticancer agent and mixtures thereof.

In certain embodiments, the compounds of the present invention can be formulated in a pharmaceutical dosage form in combination with a μ -opioid agonist. μ -opioid agonists, which may be included in the formulations of the present invention include but are not limited to include alfentanil, allylprodine, alphaprodine, anileridine, benzylmorphine, bezitramide, buprenorphine, butorphanol, clonitazene, codeine, desomorphine, dextromoramide, dezocine, diampromide, diamorphine, dihydrocodeine, dihydromorphine, dimenoxadol, dimepheptanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, eptazocine, ethoheptazine, ethylmethylthiambutene, ethylmorphine, etonitazene fentanyl, heroin, hydrocodone, hydromorphone, hydroxypethidine, isomethadone, ketobemidone, levorphanol, levophenacymorphan, lofentanil, meperidine, meptazinol, metazocine, methadone, metopon, morphine, myrophine, nalbuphine, narceine, nicomorphine, norlevorphanol, normethadone, nalorphine, normorphine, norpipanone, opium, oxycodone, oxymorphone, papaveretum, pentazocine, phenadoxone, phenomorphan, phenazocine, phenoperidine, piminodine, piritramide, proheptazine, promedol, properidine, propiram, propoxyphene, sufentanil, tilidine, tramadol, pharmaceutically acceptable salts thereof, and mixtures thereof.

In certain preferred embodiments, the μ -opioid agonist is selected from codeine, hydromorphone, hydrocodone, oxycodone, dihydrocodeine, dihydromorphine, morphine, tramadol, oxymorphone, pharmaceutically acceptable salts thereof, and mixtures thereof.

In another embodiment of the invention, the medicament comprises a mixture of a Cox-II inhibitor and an inhibitor of 5-lipoxygenase for the treatment of pain and/or inflammation. Suitable Cox-II inhibitors and 5-lipoxygenase inhibitors, as well as combinations thereof are described in U.S. Patent No. 6,136,839, which is hereby incorporated by reference in its entirety. Cox-II inhibitors include, but are not limited to rofecoxib (Vioxx), celecoxib (Celebrex), DUP-697, flosulide, meloxicam, 6-MNA, L-745337, nabumetone, nimesulide, NS-398, SC-5766, T-614, L-768277, GR-253035, JTE-522, RS-57067-000, SC-58125, SC-078, PD-138387, NS-398, flosulide, D-1367, SC-

5766, PD-164387, etoricoxib, valdecoxib and parecoxib or pharmaceutically acceptable salts, enantiomers or tautomers thereof.

The compounds of the present invention can also be combined in dosage forms with non-opioid analgesics, e.g., non-steroidal anti-inflammatory agents, including aspirin, ibuprofen, diclofenac, naproxen, benoxaprofen, flurbiprofen, fenoprofen, flubufen, ketoprofen, indoprofen, piroprofen, carprofen, oxaprozin, pramoprofen, muprofen, trioxaprofen, suprofen, aminoprofen, tiaprofenic acid, fluprofen, bucloxic acid, indomethacin, sulindac, tolmetin, zomepirac, tiopinac, zidometacin, acetaminophen, fentiazac, clidanac, oxpinac, mefenamic acid, meclofenamic acid, flufenamic acid, niflumic acid, tolfenamic acid, diflunisal, flufenisal, piroxicam, sudoxicam or isoxicam, pharmaceutically acceptable salts thereof, and mixtures thereof. Other suitable non-opioid analgesics which may be included in the dosage forms of the present invention include the following, non-limiting, chemical classes of analgesic, antipyretic, nonsteroidal antiinflammatory drugs: salicylic acid derivatives, including aspirin, sodium salicylate, choline magnesium trisalicylate, salsalate, diflunisal, salicylsalicylic acid, sulfasalazine, and olsalazin; para-aminophenol derivatives including acetaminophen; indole and indene acetic acids, including indomethacin, sulindac, and etodolac; heteroaryl acetic acids, including tolmetin, diclofenac, and ketorolac; anthranilic acids (fenamates), including mefenamic acid, and meclofenamic acid; enolic acids, including oxicams (piroxicam, tenoxicam), and pyrazolidinediones (phenylbutazone, oxyphenthartazone); and alkanones, including nabumetone. For a more detailed description of the NSAIDs that may be included within the medicaments employed in the present invention, see Paul A. Insel Analgesic-Antipyretic and Antiinflammatory Agents and Drugs Employed in the treatment of Gout in Goodman & Gilman's The Pharmacological Basis of Therapeutics, 617-57 (Perry B. Molinoff and Raymond W. Ruddon, Eds., Ninth Edition, 1996), and Glen R. Hanson Analgesic, Antipyretic and Anti-Inflammatory Drugs in Remington: The Science and Practice of Pharmacy Vol II, 1196-1221 (A. R. Gennaro, Ed. 19th Ed. 1995) which are hereby incorporated by reference in their entireties.

In certain embodiments, the compounds of the present invention can be formulated in a pharmaceutical dosage form in combination with antimigraine agents. Antimigraine agents include, but are not limited to, alpiropride, dihydroergotamine, dolasetron, ergocornine, ergocorninine, ergocryptine, ergot, ergotamine, flumetorexone acetate, fonazine, lisuride, lomerizine, methysergide, oxetorone, pizotyline, and mixtures

thereof.

The other therapeutic agent can also be an adjuvant to reduce any potential side effects such as, for example, an antiemetic agent. Suitable antiemetic agents include, but are not limited to, metoclopramide, domperidone, prochlorperazine, promethazine, chlorpromazine, trimethobenzamide, ondansetron, granisetron, hydroxyzine, acethylleucine monoethanolamine, alizapride, azasetron, benzquinamide, bietanautine, bromopride, buclizine, clebopride, cyclizine, dimenhydrinate, diphenidol, dolasetron, meclizine, methallatal, metopimazine, nabilone, oxyperndyl, pipamazine, scopolamine, sulphiride, tetrahydrocannabinols, thiethylperazine, thioproperazine, tropisetron, and mixtures thereof.

In certain embodiments, the compounds of the present invention can be formulated in a pharmaceutical dosage form in combination with β -adrenergic blockers. Suitable β -adrenergic blockers include, but are not limited to, acebutolol, alprenolol, amosulablol, arotinolol, atenolol, befunolol, betaxolol, bevantolol, bisoprolol, bopindolol, bucumolol, bufetolol, bufuralol, bunitrolol, bupranolol, butidrine hydrochloride, butofilolol, carazolol, carteolol, carvedilol, celiprolol, cetamolol, cloranolol, dilevalol, epanolol, esmolol, indenolol, labetalol, levobunolol, mepindolol, metipranolol, metoprolol, moprolol, nadolol, nadoxolol, nebivalol, nifenalol, nipradilol, oxprenolol, penbutolol, pindolol, practolol, pronethalol, propranolol, sotalol, sulfinalol, talinolol, tertatolol, tilisolol, timolol, toliprolol, and xibenolol.

In certain embodiments, the compounds of the present invention can be formulated in a pharmaceutical dosage form in combination with anticonvulsants. Suitable anticonvulsants include, but are not limited to, acetylpheneturide, albutoin, aloxidone, aminogluthethimide, 4-amino-3-hydroxybutyric acid, atrolactamide, beclamide, buramate, calcium bromide, carbamazepine, cinromide, clomethiazole, clonazepam, decimemide, diethadione, dimethadione, doxenitroin, eterobarb, ethadione, ethosuximide, ethotoin, felbamate, fluoresone, gabapentin, 5-hydroxytryptophan, lamotrigine, magnesium bromide, magnesium sulfate, mephentyoin, mephobarbital, metharbital, methetoin, methsuximide, 5-methyl-5-(3-phenanthryl)-hydantoin, 3-methyl-5-phenylhydantoin, narcobarbital, nimetazepam, nitrazepam, oxcarbazepine, paramethadione, phenacemide, phenetharbital, pheneturide, phenobarbital, phensuximide, phenylmethylbarbituric acid, phenytoin, phethenylate sodium, potassium bromide, pregabalin, primidone, progabide, sodium bromide, solanum, strontium bromide, suclofenide, sulthiame, tetrantoin, tiagabine, topiramate, trimethadione, valproic acid, valpromide, vigabatrin, and zonisamide.

In certain embodiments, the compounds of the present invention can be formulated in a pharmaceutical dosage form in combination with antidepressants. Suitable antidepressants include, but are not limited to, binedaline, caroxazone, citalopram, dimethazan, fencamine, indalpine, indeloxazine hydrochloride, nefopam, nomifensine, oxitriptan, oxypertine, paroxetine, sertraline, thiazesim, trazodone, benmoxine, iproclozide, iproniazid, isocarboxazid, nialamide, octamoxin, phenelzine, cotinine, rolicyprine, rolipram, maprotiline, metralindole, mianserin, mirtazepine, adinazolam, amitriptyline, amitriptylinoxide, amoxapine, butriptyline, clomipramine, demexiptiline, desipramine, dibenzepin, dimetacrine, dothiepin, doxepin, fluacizine, imipramine, imipramine N-oxide, iprindole, lofepramine, melitracen, metapramine, nortriptyline, noxiptilin, opipramol, pizotiline, propizepine, protriptyline, quinupramine, tianeptine, trimipramine, adrafinil, benactyzine, bupropion, butacetin, dioxadrol, duloxetine, etoperidone, febarbamate, femoxetine, fenpentadiol, fluoxetine, fluvoxamine, hematoporphyrin, hypericin, levophacetoperane, medifoxamine, milnacipran, minaprine, moclobemide, nefazodone, oxaflozane, piberaline, prolintane, pyrisuccideanol, ritanserin, roxindole, rubidium chloride, sulpiride, tandospirone, thozalinone, tofenacin, toloxatone, tranlycypromine, L-tryptophan, venlafaxine, viloxazine, and zimeldine.

In certain embodiments, the compounds of the present invention can be formulated in a pharmaceutical dosage form in combination with Ca²⁺-channel blockers. Suitable Ca²⁺-channel blockers include, but are not limited to, bepridil, clentiazem, diltiazem, fendiline, gallopamil, mibefradil, prenylamine, semotiadil, terodiline, verapamil, amlodipine, aranidipine, barnidipine, benidipine, cilnidipine, efonidipine, elgodipine, felodipine, isradipine, lacidipine, lercanidipine, manidipine, nicardipine, nifedipine, nilvadipine, nimodipine, nisoldipine, nitrendipine, cinnarizine, flunarizine, lidoflazine, lomerizine, bencyclane, etafenone, fantofarone, and perhexiline.

In certain embodiments, the compounds of the present invention can be formulated in a pharmaceutical dosage form in combination with anticancer agents. Suitable anticancer agents include, but are not limited to, acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; aminoglutethimide; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin;

carmustine; carubicin hydrochloride; carzelesin; cedefingol; chlorambucil; cirolemycin;
 cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine;
 dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine;
 dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride;
 droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate;
 eflornithine hydrochloride; elsamitrucin; enloplatin; enpromate; epipropidine; epirubicin
 hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate
 sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride;
 fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurcitabine;
 fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea;
 idarubicin hydrochloride; ifosfamide; ilmofofosine; interleukin II (including recombinant
 interleukin II, or rIL2), interferon alfa-2a; interferon alfa-2b; interferon alfa-n1 ; interferon
 alfa-n3; interferon beta-I a; interferon gamma-I b; iproplatin; irinotecan hydrochloride;
 lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol
 sodium; lomustine; losoxantrone hydrochloride; masoprocil; maytansine; mechlorethamine
 hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril;
 mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedapa;
 mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper;
 mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin;
 ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin
 sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin;
 plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride;
 puromycin; puromycin hydrochloride; pyrazofurin; riboprine; rogletimide; safingol;
 safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin;
 spirogermanium hydrochloride; spiromustine; spiroplatin; streptonigrin; streptozocin;
 sulofenur; talisomycin; tecogalan sodium; tegafur; teloxantrone hydrochloride; temoporfin;
 teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin;
 tirapazamine; toremifene citrate; trestolone acetate; triciribine phosphate; trimetrexate;
 trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa;
 vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate;
 vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine
 sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; zorubicin hydrochloride. Other
 anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-

ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetorelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatam; cypemycin; cytarabine; ocfosfate; cytolytic factor; cytostatin; daclicximab; decitabine; dehydrodidemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane;

iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin A; marimastat; masoprocil; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; monoclonal antibody, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; multiple drug resistance gene inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII

retinamide; roglitimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safangol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction modulators; single chain antigen binding protein; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; totipotent stem cell factor; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer.

The compounds of the present invention and the other therapeutic agent can act additively or, more preferably, synergistically. In a preferred embodiment, a composition comprising a compounds of the present invention is administered concurrently with the administration of another therapeutic agent, which can be part of the same composition or in a different composition from that comprising the compounds of the present invention. In another embodiment, a composition comprising the compounds of the present invention is administered prior to or subsequent to administration of another therapeutic agent.

The compounds of the present invention when administered, e.g., via the oral, parenteral or topical routes to mammals, can be in a dosage in the range of about 0.01 mg/kg to about 3000 mg/kg body weight of the patient per day, preferably about 0.01 mg/kg to about 1000 mg/kg body weight per day administered singly or as a divided dose. However, variations will necessarily occur depending upon the weight and physical condition (e.g., hepatic and renal function) of the subject being treated, the affliction to be

treated, the severity of the symptoms, the route of administration, the frequency of the dosage interval, the presence of any deleterious side-effects, and the particular compound utilized, among other things.


The compounds of the present invention preferably have a binding affinity K_i for the human ORL-1 receptor of about 500 nM or less; 100 nM or less; 50 nM or less; 20 nM or less or 5 nM or less. The binding affinity K_i can be measured by one skilled in the art by an assay utilizing membranes from recombinant HEK-293 cells expressing the human opioid receptor-like receptor (ORL-1) as described below.

The following examples illustrate various aspects of the present invention, and are not to be construed to limit the claims in any manner whatsoever.

EXAMPLE 1

SYNTHESIS OF BENZOXAZOLONE HEAD GROUPS.

The head groups of the present invention were synthesized according to the following procedure:



Procedure:

To a mixture of **1** (1.09 g, 10 mmol), **2** (1.99 g, 10 mmol) and acetic acid (0.60 g, 10 mmol) in 50 mL of dichloroethane, was added sodium triacetoxymethylborohydride (2.97 g, 14 mmol). The mixture was stirred at room temperature overnight. The mixture was filtered through Celite and 1 N NaOH (50 mL) was added to quench the reaction. The organic layer was separated and the aqueous layer was extracted with EtOAc (2 x 30 mL). The combined organic layers were dried over K_2CO_3 , filtered and evaporated in vacuum to give crude **3** as a brown solid (2.75 g, yield: 94%).

1H NMR ($CDCl_3$): δ 1.20-1.60 (m, 11H), 2.00 (dd, 2H), 2.9 (m, 2H), 3.40 (m, 1H), 4.00 (m, 2H), 6.60-6.85 (m, 4H).

To an ice cooled solution of crude **3** (12.0 g, 40 mmol) and DIEA (20.8 mL, 120 mmol) in 200 mL of THF, was added a solution of triphosgene (4.32 g, 14.4 mmol) in 200 mL of THF. After the addition was complete the ice bath was removed and the mixture stirred at room temperature overnight. The solids were filtered off and the filtrate evaporated in vacuum. The residual brown oil was dissolved in EtOAc and washed with saturated aqueous K_2CO_3 . The organic phase was dried over K_2CO_3 , filtered and evaporated in vacuum to give a red oil which was filtered through a column of silica gel eluting with a mixture of 5% Et_3N , 25% EtOAc and 70% hexane. The selected fractions were combined and the solvent evaporated in vacuum to give a brown solid which was crystallized from EtOAc to give pure **4** (10.0 g, 78% yield).

1H NMR ($CDCl_3$): δ 1.50 (s, 9H), 1.85 (d, 2H), 2.25 (m, 2H), 2.85 (m, 2H), 4.20-4.45 (m, 3H), 7.00-7.25 (m, 4H).

A solution of **4** (4.0 g, 17.2 mmol) in 30% TFA/dichloromethane (25 mL) was stirred at room temperature for 3 h. The solvent was evaporated in vacuum and saturated aqueous K_2CO_3 was added to the oily residue. The resulting mixture was extracted with dichloromethane (3 x 50 mL). The combined organic extracts were dried over K_2CO_3 , filtered and evaporated in vacuum to give the crude product. Chromatography on silica gel eluting with a mixture of 10% Et_3N , 60% EtOAc and 30% hexane gave **5** as a yellow solid (1.82 g, 66% yield).


MS: m/z 450

1H NMR ($CDCl_3$): δ 1.75-2.10 (m, 3H), 2.30 (d, 2H), 2.80 (m, 2H), 3.20 (m, 2H), 4.25 (m, 1H), 7.00-7.25 (m, 4H).

EXAMPLE 2

ATTACHMENT OF TAIL GROUPS

Tail groups were attached to the head groups according to the following procedures:



General procedure for alkylation:

To a solution of the amine (1 eq) and triethylamine (1 eq) in dimethylformamide, was added 1 eq of alkyl bromide or chloride in one portion. The mixture was stirred and heated at 80°C over night. TLC indicated the reaction was complete. The reaction was quenched by the addition of water followed by 1 N NaOH to pH 10. The mixture was extracted 2x with Et₂O. The combined organic extracts were dried over potassium carbonate and the solvent evaporated, followed by chromatography to give the pure product.

General procedure for reductive amination:

To a mixture of ketone or aldehyde (1 eq), amine (1 eq), and acetic acid (1 eq) in methanol, was added sodium cyanoborohydride (1.4 eq) in one portion. The mixture was stirred over night at room temperature. TLC indicated the reaction was complete. The reaction was quenched by the addition of water followed by 1 N NaOH to pH 10. The mixture was extracted 2x with Et₂O. The combined organic extracts were dried over potassium carbonate and the solvent evaporated, followed by chromatography to give the pure product.

The following compounds were prepared by attaching the tail groups using the general procedures described:

3-[1-(naphth-2-yl-methyl)-4-piperidiny]-2H-benzoxazol-2-one

3-[1-(naphth-1-yl-methyl)-4-piperidiny]-2H-benzoxazol-2-one

3-[1-(*p*-phenylbenzyl)-4-piperidiny]-2H-benzoxazol-2-one

3-[1-(*p*-benzyloxybenzyl)-4-piperidiny]-2H-benzoxazol-2-one

3-[1-(*p*-cyanobenzyl)-4-piperidiny]-2H-benzoxazol-2-one

MS: m/z 334.4 (M+1)

3-[1-(3,3-diphenylpropyl)-4-piperidiny]-2H-benzoxazol-2-one

3-[1-[4,4-Bis-(4-fluorophenyl)butyl]-4-piperidiny]-2H-benzoxazol-2-one

MS: m/z 463.6 (M+1).

3-[1-(2-phenylethyl)-4-piperidiny]-2H-benzoxazol-2-one

3-[1-(cyclooctylmethyl)-4-piperidiny]-2H-benzoxazol-2-one

LC: 100%

MS: m/z 343.6 (M+1).

¹H-NMR (CDCl₃): δ 1.25 (m, 2H), 1.40-1.7 (m, 17H), 2.10 (m, 4H), 3.10 (m, 2H), 4.20 (m, 1H), 7.10-7.20 (4H).

¹³C-NMR (CDCl₃): δ 26.02, 26.87, 27.55, 29.27, 31.23, 35.31, 53.39, 53.70, 66.28, 110.45, 110.51, 122.45, 123.96, 130.45, 143.08, 154.51.

3-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidiny]-2H-benzoxazol-2-one

LC: 100%

MS: 349.6 (M+1)

¹H-NMR (CDCl₃): δ 1.70 (m, 1H), 2.00 (b, 2H), 2.10 (b, 1H), 2.40 (m, 4H), 2.90 (m, 5H), 3.10 (m, 2H), 4.20 (m, 1H), 7.10-7.30 (m, 8H).

3-[1-(5-methylhex-2-yl)-4-piperidiny]-2H-benzoxazol-2-one

LC: 100%

MS: 317.4 (M+1)

¹H-NMR (CDCl₃): d 0.90 (d, 6H), 1.00 (d, 3H), 1.20 (m, 3H), 1.50-1.60 (m, 4H), 1.80 (m, 2H), 2.20-2.60 (m, 5H), 2.90 (b, 2H), 4.2 (m, 1H), 6.90-7.30 (m, 4H).

3-[1-(10,11-Dihydro-5H-dibenzo[a,d]-cyclohepten-5-yl)-4-piperidinyl]-2H-benzoxazol-2-one

LC: 96.4%

¹H-NMR (CDCl₃): d 1.80 (dd, 2H), 2.00 (dt, 2H), 2.30 (dq, 2H), 2.80-2.95 (m, 4H), 4.01 (s, 1H), 4.05-4.22 (m, 3H), 7.05-7.25 (m, 12H).

3-[1-(4-propyl-cyclohexyl)-4-piperidinyl]-2H-benzoxazol-2-one

MS: m/z 343.0

3-[1-(norbornan-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one

LC: 97%

MS: m/z 313.41 (M+1)

¹H-NMR (CDCl₃): d 0.90 (m, 1H), 1.30-2.50 (m, 17H), 3.20 (m, 2H), 4.3 (m, 1H), 6.90-7.30 (m, 4H).

3-[1-(decahydro-2-naphthyl)-4-piperidinyl]-2H-benzoxazol-2-one

MS: m/z 355.4

3-[1-(3,3-dimethyl-1,5-dioxaspiro[5.5]undeca-9-yl)-4-piperidinyl]-2H-benzoxazol-2-one

MS: m/z 401.3

3-[1-[4-(1-methylethyl)-cyclohexyl]-4-piperidinyl]-2H-benzoxazol-2-one

MS: m/z 343.0

3-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one

LC: 100%

MS: m/z 335.4 (M+1)

¹H-NMR (CDCl₃): d 1.90 (m, 1H), 2.40 (m, 2H), 2.50 (m, 2H), 2.90 (m, 2H), 3.10-3.40 (m, 6H), 4.20 (m, 1H), 7.10-7.30 (m, 8H).

3-[1-(cyclooctyl)-4-piperidinyl]-2H-benzoxazol-2-one

LC: 100%

MS: m/z 329.2 (M+1)

$^1\text{H-NMR}$ (CDCl_3): δ 1.40-2.00 (m, 16H), 2.40-2.65 (m, 4H), 2.80 (m, 1H), 3.05 (m, 2H), 4.25 (m, 1H), 7.10-7.40 (m, 4H).

Other compounds within the scope of formula (I) or (IA) of the present invention can be synthesized by analogous techniques.

EXAMPLE 3

Nociceptin affinity at the ORL1 receptor for preferred compounds was obtained using the following assay:


Membranes from recombinant HEK-293 cells expressing the human opioid receptor-like receptor (ORL-1) (Receptor Biology) were prepared by lysing cells in ice-cold hypotonic buffer (2.5 mM MgCl_2 , 50 mM HEPES, pH 7.4) (10 ml/10 cm dish) followed by homogenization with a tissue grinder/teflon pestle. Membranes were collected by centrifugation at 30,000 x g for 15 min at 4°C and pellets resuspended in hypotonic buffer to a final concentration of 1-3 mg/ml. Protein concentrations were determined using the BioRad protein assay reagent with bovine serum albumen as standard. Aliquots of the ORL-1 receptor membranes were stored at -80°C.

Functional SGTPgS binding assays were conducted as follows. ORL-1 membrane solution was prepared by sequentially adding final concentrations of 0.066 mg/ml ORL-1 membrane protein, 10 mg/ml saponin, 3 mM GDP and 0.20 nM [^{35}S]GTPgS to binding buffer (100 mM NaCl, 10 mM MgCl_2 , 20 mM HEPES, pH 7.4) on ice. The prepared membrane solution (190 ml/well) was transferred to 96-shallow well polypropylene plates containing 10 ml of 20x concentrated stock solutions of agonist prepared in DMSO. Plates were incubated for 30 min at room temperature with shaking. Reactions were terminated by rapid filtration onto 96-well Unifilter GF/B filter plates (Packard) using a 96-well tissue harvester (Brandel) and followed by three filtration washes with 200 ml ice-cold binding buffer (10 mM NaH_2PO_4 , 10 mM Na_2HPO_4 , pH 7.4). Filter plates were subsequently dried at 50°C for 2-3 hours. Fifty ml/well scintillation cocktail (BetaScint; Wallac) was added and plates were counted in a Packard Top-Count for 1 min/well.

Data was analyzed using the curve fitting functions in GraphPad PRISM[®], v. 3.0 and the results are set forth in table 1 below:

TABLE 1	
Nociceptin Affinity	
Compound	calc K _i (nM)
3-[1-(naphth-2-yl-methyl)-4-piperidinyl]-2H-benzoxazol-2-one	3030
3-[1-(naphth-1-yl-methyl)-4-piperidinyl]-2H-benzoxazol-2-one	370
3-[1-(<i>p</i> -phenylbenzyl)-4-piperidinyl]-2H-benzoxazol-2-one	>10,000
3-[1-(<i>p</i> -benzyloxybenzyl)-4-piperidinyl]-2H-benzoxazol-2-one	2173
3-[1-(<i>p</i> -cyanobenzyl)-4-piperidinyl]-2H-benzoxazol-2-one	>10,000
3-[1-(3,3-diphenylpropyl)-4-piperidinyl]-2H-benzoxazol-2-one	726
3-[1-[4,4-Bis-(4-fluorophenyl)butyl]-4-piperidinyl]-2H-benzoxazol-2-one	3070
3-[1-(2-phenylethyl)-4-piperidinyl]-2H-benzoxazol-2-one	7087
3-[1-(cyclooctylmethyl)-4-piperidinyl]-2H-benzoxazol-2-one	64
3-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-2H-benzoxazol-2-one	93
3-[1-(5-methylhex-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one	60
3-[1-(10,11-Dihydro-5H-dibenzo[a,d]-cyclohepten-5-yl)-4-piperidinyl]-2H-benzoxazol-2-one	>10,000
3-[1-(3,3-dimethyl-1,5-dioxaspiro[5.5]undeca-9-yl)-4-piperidinyl]-2H-benzoxazol-2-one	>10,000
3-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one	512
3-[1-(cyclooctyl)-4-piperidinyl]-2H-benzoxazol-2-one	16

EXAMPLE 4
SYNTHESIS OF SUBSTITUTED INDOLE HEAD GROUPS



Procedure:

To a mixture of 2 (23.3 g, 0.25 mol), 1 (47.3 g, 0.25 mol), acetic acid (15 g, 0.25 mol) and molecular sieves (15 g) in 500 mL of dichloroethane, sodium triacetoxyborohydride (74.2 g, 0.35 mol) was added in one portion and the mixture stirred overnight. The molecular sieves were filtered off and 1 N NaOH (500 mL) was added to quench the reaction. The

organic layer was separated and the aqueous layer extracted with EtOAc (2 x 300 mL). The combined organic extracts were dried over K_2CO_3 , filtered and the solvent evaporated under vacuum to give crude 3 as a brown solid which was directly used in next step.

Compound 3

1H -NMR ($CDCl_3$): δ 1.50 (m, 2H), 2.05 (m, 2H), 2.20 (bt, 2H), 2.85 (m, 2H), 3.30 (m, 1H), 3.52 (s, 2H), 6.60 (d, 2H), 6.70 (t, 1H), 7.20 (m, 2H), 7.25-7.40 (m, 5H).

To an ice cooled solution of crude 3 (0.25 mol, 100% yield assumed) and DIEA (48.4 g, 0.38 mol) in 500 mL of dichloromethane, was added dropwise chloroacetyl chloride (42.4 g, 0.375 mol). After the addition was complete the ice bath was removed and the reaction mixture stirred overnight. The solvent was removed in vacuum and the residue dissolved in dichloromethane. The organic phase was washed with saturated aqueous K_2CO_3 , dried over K_2CO_3 , filtered and the solvent removed in vacuum to give a brown gum which was filtered through a column of silica gel eluting with a mixture of 10% Et_3N , 40% EtOAc and 50% hexane. The selected fractions were combined and the solvent evaporated in vacuum to give a brown solid which was further crystallized from EtOAc to give 42.2 g of 4 (49.2%, 2 steps).

Compound 4

1H NMR (DMSO): δ 1.22 (m, 2H), 1.70 (b, 2H), 2.00 (t, 2H), 2.80 (b, 2H), 3.40 (s, 2H), 3.80 (s, 2H), 4.40 (m, 1H), 7.15-7.30 (m, 7H), 7.45 (m, 3H).

A mixture of 4 (42.2 g, 0.12 mol) and $AlCl_3$ (49.2 g, 0.369 mol) was mixed in a flask by rapid stirring. The mixture was then heated in an oil bath at 130 °C. Within a few minutes the solids melted and became a dark liquid with concomitant gas evolution. After heating for 1 h the reaction mixture was cooled somewhat and while still mobile poured into a beaker containing 500 mL of ice water. The solution was basified and extracted with dichloromethane. The organic layer was dried over Na_2SO_4 , filtered and the solvent evaporated in vacuum to give a dark oil which was filtered through a column of silica gel eluting with a mixture of 10% Et_3N , 40% EtOAc and 50% hexane. The selected fractions were combined and the solvent evaporated in vacuum to give 5 as a red oil which set to a pale solid (22.0 g, 58.5%).

Compound 5

¹H NMR (CDCl₃): d 1.70 (m, 2H), 2.17 (m 2H), 2.50 (m, 2H), 3.05 (m, 2H), 3.55 (s, 2H), 3.60 (s, 2H), 4.33 (m, 1H), 7.00-7.40 (m, 9H).

To a solution of 5 (16.0 g, 0.052 mol) in 35 mL of methanol was added Pd(OH)₂ (4.0 g). The resulting suspension was hydrogenated at 50 psi for 12 h at room temperature. The solution was filtered through a pad of Celite and the pad washed with methanol (2 x 20 mL). Evaporation of the solvent in vacuum gave 6 as a pale solid (11.2 g, 100%).

Compound 6

LC: 100%

MS: m/z 217 (M+1).

¹H NMR (CDCl₃): d 1.75 (m, 3H), 2.35 (m 2H), 2.75 (m, 2H), 3.25 (m, 2H), 3.50 (s, 2H), 4.33 (m, 1H), 7.00-7.30 (m, 4H)

To a solution of 6 (8.0 g, 37.0 mmol) in 50 mL of dichloromethane was added Et₃N (4.07 g, 40.7 mmol) and BOC anhydride (8.87 g, 40.7 mmol). After stirring for 3 h saturated aqueous K₂CO₃ was added and the layers separated. The aqueous phase was extracted with dichloromethane (2 x 50 mL). The combined organic phase was dried over K₂CO₃, filtered and evaporated in vacuum to give a brown oil which was filtered through a column of silica gel eluting with a mixture of 10% Et₃N, 40% EtOAc and 50% hexane. The selected fractions were combined and the solvent evaporated in vacuum to give 7 as an off white solid (8.50 g, 73%).

Compound 7

¹H NMR (CDCl₃): d 1.50 (m, 9H), 1.70 (m 2H), 2.20-2.50 (m, 2H), 2.80-3.00 (m, 2H), 3.50 (s, 2H), 4.20-4.50 (m, 3H), 6.90-7.60 (m, 5H).

To a mixture of 7 (6.0 g, 19.0 mmol) and sodium acetate (2.58 g, 19.0 mmol) in 150 mL of methanol was added acetaldehyde (1.67 g, 38.0 mmol). The mixture was refluxed for 2 h. The solvent was evaporated in vacuum to give a dark oil which was filtered through a column of silica gel eluting with a mixture of 10% Et₃N, 40% EtOAc and 50% hexane. The

selected fractions were combined and the solvent evaporated in vacuum to give 8 as a red oil (5.90 g, 91%).

Compound 8

LC: 2 isomers in a ratio of 2:1.

¹H NMR (CDCl₃): (mixture of 2 isomers) δ 1.50 (m, 9H), 1.70 (m 2H), 2.20-2.50 (m, 6H), 2.60-3.00 (m, 2H), 4.20-4.50 (m, 3H), 6.90-7.60 (m, 5H).

A solution of 8 (5.90 g, 17.2 mmol) in 30% TFA/dichloromethane (100 mL) was stirred at room temperature for 3 h. The solvent was evaporated in vacuum and saturated aqueous K₂CO₃ was added to the oily residue. The resulting mixture was extracted with dichloromethane (3 x 150 mL). The combined organic extracts were dried over K₂CO₃, filtered and evaporated in vacuum to give the crude product. Chromatography on silica gel eluting with a mixture of 10% Et₃N, 50% EtOAc and 40% hexane gave 9 (E/Z isomers) as a yellow foam (3.60 g, 82%).


Compound 9

LC: 2 isomers in a ratio of 2:1.

MS: m/z 243.1 (M+1).

¹H NMR (CDCl₃): (mixture of 2 isomers) δ 0.85 (m, 1H), 1.50-2.00 (m, 4H), 2.20-2.50 (m, 5H), 2.60 (m, 1H), 3.10-3.50 (m, 2H), 4.30 (m, 1H), 6.90-7.60 (m, 5H).

EXAMPLE 5



To a mixture of **5** (5.50 g, 18 mmol) and sodium acetate (2.45 g, 18 mmol) in 150 mL of methanol was added acetaldehyde (1.58 g, 36 mmol). The mixture was refluxed for 2 h. The solvent was evaporated in vacuum to give a dark oil which was filtered through a column of silica gel eluting with a mixture of 10% Et₃N, 40% EtOAc and 50% hexane. The selected fractions were combined and the solvent evaporated in vacuum to give **10** as a red oil (5.90 g, 98%).

Compound 10

LC: 2 isomers in a ratio of 2:1.

MS: m/z 333.2 (M+1).

¹H NMR (CDCl₃): δ 1.70 (m, 2H), 2.17 (m, 2H), 2.30 (d, 3H), 2.50 (m, 2H), 3.05 (m, 2H), 3.55 (s, 2H), 4.33 (m, 1H), 7.00-7.40 (m, 9H), 7.6 (d, 1H).

To a solution of 10 (5.90 g, 17.7 mmol) in 30 mL of methanol was added Pd(OH)₂ (3.0 g). The resulting suspension was hydrogenated at 50 psi for 12 h at room temperature. The solution was filtered through a pad of Celite and the pad washed with methanol (2 x 20 mL). Evaporation of the solvent in vacuum gave a pale solid which was purified by chromatography on silica gel eluting with a mixture of 10% methanol and 90% EtOAc to give 11 as an off white solid (2.02 g, 50%).


Compound 11

LC: 97%

MS: m/z 245.2 (M+1)

^1H NMR (CDCl_3): d 0.85 (t, 3H), 1.26 (m, 2H), 2.00 (m, 2H), 2.43 (m, 2H), 2.90 (m, 2H), 3.3 (m, 2H), 3.4 (m, 1H), 4.4 (m, 1H), 7.05 (m, 1H), 7.15-7.30 (m, 3H).

EXAMPLE 6



Procedure:

In a manner similar to the preparation of **6**, compound **16** was prepared.

Compound **13**

LC: 89.4%

MS: m/z 281.2 ($M+1$)

^1H -NMR (mixture of trans and cis) (CDCl_3): d 0.95 (m, 3H), 1.50-2.75 (m, 5H), 2.80-3.20 (m, 1H), 3.50 (m, 2H), 3.60 (minor)+3.70 (major) (two s, 2H), 6.55-6.80 (m, 2H), 7.05-7.45 (m, 8H).

Compound **14**

MS: m/z 357.2 ($M+1$)

¹H-NMR (mixture of trans and cis) (CDCl₃): d 1.10 (m, 3H), 1.40-4.20 (m, 11H), 4.40 (m, 1H), 7.05-7.50 (m, 10H).

Compound 15

LC: 90.0%

MS: m/z 321.2 (M+1)


¹H-NMR (CDCl₃): d 1.20 (d, 3H), 1.75 (m, 1H), 2.10 (dt, 1H), 2.25 (b, 1H), 2.30 (dd, 1H), 2.75 (dd, 1H), 3.05 (m, 1H), 3.20 (m, 1H), 3.50 (m, 4H), 4.10 (m, 1H), 6.99 (m, 2H), 7.23 (m, 3H), 7.37 (m, 4H).

Compound 16

LC: 92.5%

MS: m/z 231.2 (M+1)

¹H-NMR (CDCl₃): d 1.20 (d, 3H), 1.75 (m, 1H), 2.10 (dt, 1H), 2.25 (b, 1H), 2.30 (dd, 1H), 2.75 (dd, 1H), 3.05 (m, 1H), 3.20 (m, 1H), 3.50 (m, 2H), 4.10 (m, 1H), 6.99 (m, 2H), 7.23 (m, 3H), 7.37 (m, 4H).

EXAMPLE 7

Procedure:

In a manner similar to the preparation of 11, compound 18 was prepared.

Compound 17

MS: m/z 347.3 (M+1)

Compound 18


LC: 82.6%

MS: m/z 259.3 (M+1)

¹H-NMR (CDCl₃): δ 0.80 (t, 3H), 1.20 (d, 3H), 2.00 (m, 2H), 2.30 (m, 1H), 2.65 (m, 1H), 2.82 (m, 1H), 3.15-3.25 (m, 1H), 3.32 (m, 1H), 3.45 (m, 1H), 3.65 (m, 1H), 3.75 (m, 1H), 4.25 (m, 1H), 6.90 (d, 1H), 7.05 (t, 1H), 7.25 (m, 2H).

EXAMPLE 8
ATTACHMENT OF TAIL GROUPS

Tail groups were attached to the head groups according to the following procedures:



General procedure for alkylation:

To a solution of the amine (1 eq) and triethylamine (1 eq) in dimethylformamide, was added 1 eq of alkyl bromide or chloride in one portion. The mixture was stirred and heated at 80°C over night. TLC indicated the reaction was complete. The reaction was quenched by the addition of water followed by 1 N NaOH to pH 10. The mixture was extracted 2x with Et₂O. The combined organic extracts were dried over potassium carbonate and the solvent evaporated, followed by chromatography to give the pure product.

General procedure for reductive amination:

To a mixture of ketone or aldehyde (1 eq), amine (1 eq), and acetic acid (1 eq) in methanol, was added sodium cyanoborohydride (1.4 eq) in one portion. The mixture was stirred over night at room temperature. TLC indicated the reaction was complete. The reaction was quenched by the addition of water followed by 1 N NaOH to pH 10. The mixture was extracted 2x with Et₂O. The combined organic extracts were dried over potassium carbonate and the solvent evaporated, followed by chromatography to give the pure product.

The following compounds were prepared by attaching the tail groups using the general procedures described:

1-[1-(naphth-1-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 357.2 (M+1).

1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 357.3 (M+1).

1-[1-(*p*-phenylbenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 383.2 (M+1).

1-[1-(3,3-Bis(phenyl)propyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 98.7%

MS: m/z 411.2 (M+1)

¹H-NMR (CDCl₃): d 1.65 (bd, 2H), 2.05 (bt, 2H), 2.30 (m, 4H), 2.45 (m, 2H), 3.02 (bd, 2H), 3.50 (s, 2H), 4.01 (t, 1H), 4.30 (m, 1H), 7.00 (t, 1H), 7.15-7.35 (m, 13H).

1-[1-(*p*-cyanobenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 332.2 (M+1).

1-[1-(*p*-benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 413.3 (M+1)

1-[1-(1,2,3,4-tetrahydronaphth-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 100%

MS: m/z 347.5 (M+1).

¹H NMR (CDCl₃): d 1.70 (m, 3H), 2.10 (m, 1H), 2.40 (m, 4H), 2.90 – 3.00 (m, 5H), 3.10 (m, 2H), 3.60 (s, 2H), 4.3 (m, 1H), 7.00-7.30 (m, 8H).

1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 100%

MS: m/z 315.4 (M+1).

¹H NMR (CDCl₃): d 0.90 (m, 6H), 1.00 (m, 3H), 1.20 (m, 3H), 1.5-1.8 (m, 2H), 2.2-2.6 (m, 5H), 2.90 (m, 2H), 3.60 (s, 2H), 4.2 (m, 1H), 6.90-7.30 (m, 4H).

1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 97%

MS: m/z 311.41 (M+1).

¹H NMR (CDCl₃): d 0.90 (m, 1H), 1.30-2.00 (m, 7H), 2.10-2.30 (m, 5H), 3.20 (m, 2H), 3.60 (s, 2H), 4.3 (m, 1H), 6.90-7.30 (m, 4H).

1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 100%

MS: m/z 332.4 (M+1).

¹H NMR (CDCl₃): d 1.80 (m, 2H), 2.40 (m, 2H), 2.50 (m, 2H), 2.90 (m, 2H), 3.10-3.40 (m, 5H), 3.60 (s, 2H), 4.20 (m, 1H), 7.10-7.30 (m, 8H).

1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 97%

MS: m/z 341.50 (M+1).

¹H NMR (CDCl₃): d 1.25 (m, 3H), 1.4-1.7 (m, 14H), 2.10 (m, 4H), 2.50 (m, 2H), 3.10 (m, 2H), 3.60 (s, 2H), 4.3 (m, 1H), 7.10-7.20 (m, 4H).

¹³C-NMR (CDCl₃): d 23.07, 26.04, 26.89, 27.56, 28.63, 31.27, 32.00, 35.30, 36.33, 46.63, 50.65, 54.06, 66.47, 110.90, 122.17, 124.90, 125.26, 127.94, 144.25, 175.31.

3-ethyl-1-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 375.3 (M+1).

3-ethyl-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 369.2 (M+1).

3-ethyl-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 100%

MS: m/z 342.4 (M+1).

¹H NMR (CDCl₃): d 0.80 (t, 3H), 0.90 (m, 6H), 1.00 (m, 3H), 1.20 (m, 3H), 1.5-1.8 (m, 2H), 2.2-2.6 (m, 5H), 2.90 (m, 2H), 3.40 (m, 1H), 4.3 (m, 1H), 6.90-7.30 (m, 4H).

3-ethyl-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 100%

MS: m/z 339.41 (M+1).

¹H NMR (CDCl₃): d 0.80 (m, 3H), 0.90 (m, 1H), 1.30-1.45 (m, 5H), 1.50-2.05 (m, 8H), 2.10 (m, 1H), 2.20 (m, 2H), 2.50 (m, 2H), 3.10 (m, 2H), 3.40 (m, 1H), 4.3 (m, 1H), 6.90-7.30 (m, 4H).

3-ethyl-1-[1-(decahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 381.3 (M+1).

3-ethyl-1-[1-[4-(1-methylethyl)-cyclohexyl]-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 369.3 (M+1)

¹H-NMR (CDCl₃): d 0.88 (t, 3H), 0.92 (d, 6H), 1.17 (m, 1H), 1.40 (m, 2H), 1.50-1.70 (m, 9H), 2.05 (m, 2H), 2.25 (m, 2H), 2.32-2.55 (m, 3H), 3.15 (b, 2H), 3.43 (t, 1H), 4.35 (m, 1H), 7.05 (t, 1H), 7.22 (d, 1H), 7.28 (m, 2H).

3-ethyl-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 361.2 (M+1).

3-ethyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 97%

MS: m/z 369.50 (M+1).

¹H NMR (CDCl₃): d 0.80 (t, 3H), 1.25 (m, 3H), 1.4-1.7 (m, 14H), 2.10 (m, 6H), 2.50 (m, 2H), 3.10 (m, 2H), 3.40 (m, 1H), 4.3 (m, 1H), 7.10-7.20 (m, 4H).

3-ethylidene-1-[1-(benzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 333.2 (M+1)

¹H-NMR (CDCl₃): d 1.70 (m, 2H), 2.15 (dt, 2H), 2.28 (d, 3H), 2.47 (m, 2H), 3.05 (b, 2H), 3.57 (s, 2H), 4.34 (m, 1H), 7.02 (t, 1H), 7.08-7.40 (m, 8H), 7.58 (d, 1H).

3-ethylidene-1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 405.2

3-ethylidene-1-[1-(3,3-diphenylpropyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: >97% (2 isomers combined).

MS: m/z 437.5 (M+1).

¹H NMR (CDCl₃): d 1.70-1.80 (m, 3H), 2.10 (m, 2H), 2.20-2.40 (m, 8H), 3.10 (m, 2H), 4.10 (m, 1H), 4.3 (m, 1H), 7.00-7.30 (m, 15H).

3-ethylidene-1-[1-(*p*-cyanobenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: >97% (2 isomers combined).

MS: m/z 358.5 (M+1).

^1H NMR (CDCl_3): δ 1.80 (m, 4H), 2.10–2.60 (m, 5H), 3.10 (m, 2H), 3.70 (s, 2H), 4.3 (m, 1H), 6.90–7.60 (m, 8H).

3-ethylidene-1-[1-(*p*-benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

MS: m/z 405.2.

3-ethylidene-1-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: >97% (2 isomers combined).

MS: m/z 373.5 (M+1).

^1H NMR (CDCl_3): δ 1.70–3.10 (m, 18H), 4.3 (m, 1H), 7.00–7.30 (m, 9H).

3-ethylidene-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: >97% (2 isomers combined).

MS: m/z 367.5 (M+1).

^1H NMR (CDCl_3): δ 0.90 (m, 1H), 1.30–2.00 (m, 7H), 2.10–2.30 (m, 5H), 3.20 (m, 2H), 3.60 (s, 2H), 4.3 (m, 1H), 6.90–7.30 (m, 5H).

3-ethylidene-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: >97% (2 isomers combined).

MS: m/z 341.4 (M+1).

^1H NMR (CDCl_3): δ 0.90–2.6 (m, 24H), 2.90 (m, 2H), 4.2 (m, 1H), 6.90–7.30 (m, 5H).

3-ethylidene-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: >97% (2 isomers combined).

MS: m/z 337.41 (M+1).

^1H NMR (CDCl_3): δ 0.90 (m, 1H), 1.30–2.50 (m, 17H), 3.10 (m, 2H), 4.3 (m, 1H), 6.90–7.30 (m, 5H).

3-ethylidene-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: >97% (2 isomers combined).

MS: m/z 359.4 (M+1).

¹H NMR (CDCl₃): d 1.80 - 3.10 (m, 17H), 4.20 (m, 1H), 7.10-7.30 (m, 9H).

3-ethylidene-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: >97% (2 isomers combined).

MS: m/z 367.50 (M+1).

¹H NMR (CDCl₃): d 1.25 (m, 3H), 1.4-1.7 (m, 21H), 2.10 - 2.50 (m, 2H), 3.10 (m, 2H), 4.3 (m, 1H), 6.90-7.60 (m, 5H).

1-[1-(3,3-Bis(phenyl)propyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 100%

MS: m/z 425.3 (M+1)

¹H-NMR (CDCl₃): d 1.20 (d, 3H), 1.69 (bd, 1H), 1.95 (dt, 1H), 2.13-2.30 (m, 5H), 2.72 (bd, 1H), 2.98 (bd, 1H), 3.15 (dq, 1H), 3.50 (s, 2H), 4.03 (dt, 1H), 4.12 (t, 1H), 6.94 (d, 1H), 7.00 (t, 1H), 7.10-7.30 (m, 12H).

1-[1-(benzyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 100%

MS: m/z 321.2 (M+1)

¹H-NMR (CDCl₃): d 1.20 (d, 3H), 1.70 (m, 1H), 2.10 (dt, 1H), 2.23 (m, 1H), 2.35 (dd, 1H), 2.78 (d, 1H), 3.05 (m, 1H), 3.20 (dq, 1H), 3.51 (m, 4H), 4.10 (dt, 1H), 7.00 (m, 2H), 7.25 (m, 3H), 7.38 (m, 4H).

1-[1-(4-propyl-cyclohexyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 96.2%

MS: m/z 355.2 (M+1)

¹H-NMR (CDCl₃): d 0.85 (m, 3H), 1.15 (m, 3H), 1.22-1.85 (m, 13H), 2.05-2.90 (m, 6H), 2.95-3.20 (m, 2H), 3.50 (s, 2H), 4.05 (m, 1H), 7.00 (m, 2H), 7.22 (m, 2H).

1-[1-(5-methylhex-2-yl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 100%

MS: m/z 329.2 (M+1)

¹H-NMR (CDCl₃): δ 0.85 (m, 9H), 1.15 (m, 3H), 1.20-1.75 (m, 6H), 2.25 (m, 1H), 2.45-2.75 (m, 4H), 2.88 (m, 1H), 3.10 (m, 1H), 3.50 (s, 2H), 4.05 (m, 1H), 6.98 (m, 2H), 7.25 (m, 2H).

1-[1-(decahydro-2-naphthyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 95.3%

MS: m/z 367.2 (M+1)

¹H-NMR (CDCl₃): δ 1.11 (d, 3H), 1.16-1.85 (m, 16H), 2.20 (m, 1H), 2.35 (m, 2H), 2.52 (m, 2H), 2.75 (m, 1H), 3.02 (m, 2H), 3.50 (s, 2H), 4.05 (m, 1H), 6.96 (m, 2H), 7.20 (m, 2H).

1-[1-(4-(1-methylethyl)-cyclohexyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 96.1%

MS: m/z 355.2 (M+1)

¹H-NMR (CDCl₃): δ 0.80 (m, 6H), 1.15 (m, 3H), 1.22-1.48 (m, 3H), 1.50-1.90 (m, 6H), 2.15-2.90 (m, 4H), 2.95-3.25 (m, 2H), 3.50 (s, 2H), 4.10 (m, 1H), 6.95 (m, 2H), 7.22 (m, 2H).

1-[1-(cyclooctylmethyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 100%

MS: m/z 355.2 (M+1)

¹H-NMR (CDCl₃): δ 1.12 (d, 3H), 1.15-1.75 (m, 16H), 1.92-2.10 (m, 3H), 2.20 (m, 2H), 2.73 (m, 1H), 3.00 (m, 1H), 3.12 (dq, 1H), 3.50 (s, 2H), 4.05 (dt, 1H), 6.99 (m, 2H), 7.20 (m, 2H).

3-ethyl-1-[1-(3,3-Bis(phenyl)propyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 96.3%

MS: m/z 453.3 (M+1)

¹H-NMR (CDCl₃): δ (two t, 3H), 1.18 (d, 3H), 1.70 (m, 1H), 1.90-2.05 (m, 3H), 2.12-2.30 (m, 5H), 7.73 (m, 1H), 2.97 (bd, 1H), 3.10-3.30 (m, 1H), 3.38 (t, 1H), 3.90-4.05 (m, 1H), 4.12 (q, 1H), 6.90-7.00 (two d, 1H), 7.02 (t, 1H), 7.12-7.32 (m, 12H).

3-ethyl-1-[1-(4-propylcyclohexyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 93.2%

MS: m/z 383.3 ($M+1$)

$^1\text{H-NMR}$ (CDCl_3): δ 0.75-0.95 (m, 6H), 1.05-1.20 (m, 5H), 1.20-1.35 (m, 4H), 1.35-1.75 (m, 6H), 1.75-1.90 (m, 2H), 1.95-2.05 (m, 2H), 2.15-2.45 (m, 3H), 2.55 (d, 0.5H), 2.75 (d, 0.5H), 2.95-3.15 (m, 2H), 3.38 (t, 1H), 3.90-4.10 (m, 1H), 6.90-7.05 (2H), 7.20-7.25 (m, 2H).

3-ethyl-1-[1-(5-methylhex-2-yl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

LC: 92.3%

MS: m/z 357.4 ($M+1$)

$^1\text{H-NMR}$ (CDCl_3): δ 0.75-0.95 (m, 10H), 1.10 (d, 3H), 1.15-1.40 (m, 3H), 1.40-1.75 (m, 4H), 1.97-2.10 (m, 2H), 2.20 (m, 1H), 2.43-2.75 (m, 4H), 2.80-2.95 (m, 1H), 3.00-3.25 (m, 1H), 3.40 (t, 1H), 3.90-4.10 (m, 1H), 6.90-7.05 (m, 2H), 7.25 (m, 2H).

3-ethyl-1-[1-[4-(1-methylethyl)cyclohexyl]-3-methyl-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 94.7%

MS: m/z 383.4 ($M+1$)

$^1\text{H-NMR}$ (CDCl_3): δ 0.75-1.05 (m, 8H), 1.10-1.50 (m, 7H), 1.50-1.90 (m, 7H), 1.90-2.10 (m, 2H), 2.15-2.43 (m, 3H), 2.55 (d, 0.5H), 2.75 (d, 0.5H), 2.90-3.25 (m, 3H), 3.40 (t, 1H), 3.90-4.10 (m, 1H), 6.90-7.01 (m, 2H), 7.25 (m, 2H).

3-ethyl-1-[1-(decahydro-2-naphthyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one

LC: 94.3%

MS: m/z 395.3 ($M+1$)

$^1\text{H-NMR}$ (CDCl_3): δ 1.75-1.90 (two t, 3H), 1.10 (d, 3H), 1.15-1.90 (m, 15H), 2.00 (m, 2H), 2.20 (bs, 1H), 2.40 (m, 2H), 2.45-2.60 (m, 2H), 2.75 (m, 1H), 2.90-3.20 (m, 2H), 3.40 (bs, 1H), 3.90-4.15 (m, 1H), 6.90-7.05 (m, 2H), 7.25 (m, 2H).

Other compounds within the scope of formula (II) or (IIA) of the present invention can be synthesized by analogous techniques.

EXAMPLE 9

Nociceptin affinity at the ORL1 receptor for preferred compounds was obtained using the following assay:

Membranes from recombinant HEK-293 cells expressing the human opioid receptor-like receptor (ORL-1) (Receptor Biology) were prepared by lysing cells in ice-cold hypotonic buffer (2.5 mM $MgCl_2$, 50 mM HEPES, pH 7.4) (10 ml/10 cm dish) followed by homogenization with a tissue grinder/teflon pestle. Membranes were collected by centrifugation at 30,000 x g for 15 min at 4°C and pellets resuspended in hypotonic buffer to a final concentration of 1-3 mg/ml. Protein concentrations were determined using the BioRad protein assay reagent with bovine serum albumen as standard. Aliquots of the ORL-1 receptor membranes were stored at -80°C.

Functional SGTPgS binding assays were conducted as follows. ORL-1 membrane solution was prepared by sequentially adding final concentrations of 0.066 mg/ml ORL-1 membrane protein, 10 mg/ml saponin, 3 mM GDP and 0.20 nM [^{35}S]GTPgS to binding buffer (100 mM NaCl, 10 mM $MgCl_2$, 20 mM HEPES, pH 7.4) on ice. The prepared membrane solution (190 ml/well) was transferred to 96-shallow well polypropylene plates containing 10 ml of 20x concentrated stock solutions of agonist prepared in DMSO. Plates were incubated for 30 min at room temperature with shaking. Reactions were terminated by rapid filtration onto 96-well Unifilter GF/B filter plates (Packard) using a 96-well tissue harvester (Brandel) and followed by three filtration washes with 200 ml ice-cold binding buffer (10 mM NaH_2PO_4 , 10 mM Na_2HPO_4 , pH 7.4). Filter plates were subsequently dried at 50°C for 2-3 hours. Fifty ml/well scintillation cocktail (BetaScint; Wallac) was added and plates were counted in a Packard Top-Count for 1 min/well.

Data was analyzed using the curve fitting functions in GraphPad PRISM®, v. 3.0 and the results are set forth in table 2 below:


TABLE 2	
Nociceptin Affinity	
Compound	calc K_i (nM)
3-ethylidene-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	11.1
3-ethylidene-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	19
3-ethylidene-1-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	16.7
3-ethylidene-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	20.7
3-ethylidene-1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	630
3-ethylidene-1-[1-(<i>p</i> -benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	516
3-ethylidene-1-[1-(benzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	1854
3-ethylidene-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	22.3
3-ethylidene-1-[1-(3,3-diphenylpropyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	100.7
3-ethylidene-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	922
3-ethylidene-1-[1-(<i>p</i> -cyanobenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	7652
3-ethyl-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	4
3-ethyl-1-[1-[4-(1-methylethyl)-cyclohexyl]-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	.86
3-ethyl-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	40
3-ethyl-1-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	124
3-ethyl-1-[1-(decahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	3.6
3-ethyl-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	43
3-ethyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	9
3-ethyl-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	82.7
1-[1-(naphth-1-ylmethyl)-4-piperidinyl]-1,3-dihydro-2H-	92

indole-2-one	
1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	107
1-[1-(<i>p</i> -phenylbenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	1362
1-[1-(3,3-Bis(phenyl)propyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	12.5
1-[1-(<i>p</i> -cyanobenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	1267
1-[1-(<i>p</i> -benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	32
1-[1-(1,2,3,4-tetrahydronaphth-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	28.7
1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	7.4
1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	215
1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	18.7
1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	54.3
1-[1-(benzyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	> 10,000
1-[1-(4-propyl-cyclohexyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	2435
1-[1-(5-methylhex-2-yl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	4335
1-[1-(decahydro-2-naphthyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	366
1-[1-(4-(1-methylethyl)-cyclohexyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	167
1-[1-(cyclooctylmethyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	189

EXAMPLE 10

SYNTHESIS OF CERTAIN HEAD GROUPS.

SCHEME 1:

Procedure:

To a mixture of 4 (21.6 g, 0.2 mole), 5 (15.6 g, 0.1 mole), acetic acid (6 g, 0.1mole) in 500 ml of dichloroethane, 29.7 g of sodium triacetoxyborohydride (0.14 mol, 1.4 eq) was added in one portion. Gas evolves between 30 min and 1 hr. The mixture was stirred over night. TLC

indicated the reaction is complete. 1 N NaOH (500 ml) was added to quench the reaction. The organic layer was separated and the aqueous layer was extracted by EtOAc (300 ml x2). The combined organics were dried over potassium carbonate and the solvent evaporated to give a red oil which was column filtrated (5%Et₃N, 25%EtOAc and 70%Hexane) to give 14 g of product 6 as a white solid (54%).

Compound 6

MS: m/z 249.3 (M+1).

¹H NMR (CDCl₃): δ 1.50 – 1.90 (m, 6H), 2.05 (m, 2H), 3.30 (m, 4H), 3.95 (s, 4H), 6.60 – 6.80 (m, 4H).

To a solution of 13.5 g of 6 (54.4 mmol) in 50 ml of acetonitrile, 11.02 g of carbonyldiimidazole was added in one portion. The mixture was stirred over night. Solid precipitated out of solution which was filtered and washed by H₂O and TBME to give 7.5 g of product. The filtrate was evaporated and the crude material was dissolved in EtOAc, washed with water and saturated potassium carbonate solution. The organics were dried over potassium carbonate. The solvent was evaporated to give a second batch of solid with a pink color which was column filtrated (10%Et₃N, 40%EtOAc and 50%Hexane) to give another 4.5 g of product 7 (81 %, combined).

Compound 7

MS: m/z 274.7 (M+1).

¹H NMR (CDCl₃): δ 1.50 – 1.90 (m, 7H), 2.50 (m, 2H), 4.00 (m, 4H), 4.50 (m, 1H), 7.10 (m, 3H), 7.25 (m, 1H).

A mixture of 7 (7.5 g, 27.4 mmole) and 8.26 g of PPTS in 50 ml of acetone and H₂O (10:1) was stirred in refluxed over night. The mixture was cooled to room temperature and acetone was evaporated. Addition of water to the mixture initiated crystalization to give 3 g of product 8 (47.4%).

Compound 8

MS: m/z 231 (M+1).

¹H NMR (CDCl₃): δ 2.20 (m, 2H), 2.60 (m, 2H), 4.50 (m, 1H), 7.10 (m, 4H), 9.5 (br, 1H).

To a mixture of 8 (7.75 g, 33.65 mmole), benzylamine (3.61 g, 33.65 mmole), acetic acid (2.0 g, 33.65 mmole) in 150 ml of dichloroethane, 10.3 g of sodium triacetoxymethylborohydride (47.1 mmol, 1.4 eq) was added in one portion. Gas evolves between 30 min and 1 hr. The mixture was stirred over night. TLC indicated the reaction was complete. 1 N NaOH (500 ml) was added to quench the reaction. The organic layer was separated and the aqueous layer was extracted with EtOAc (300 ml x2). The combined organics were dried over potassium carbonate and the solvent was evaporated to give a brown solid, which was column filtrated (5%Et3N, 25%EtOAc and 70%Hexane to 10%Et3N, 40%EtOAc and 50%Hexane) to give 4.7 g of product 10 as a white solid (53.4%) and 3.01 g of product 9 as a white solid (34.2%).

Compound 9

MS: m/z 322 (M+1).

¹H NMR (CDCl₃): d 1.40 (m, 2H), 1.80-2.35 (m, 6H), 2.70 (m, 1H), 3.86 (s, 2H), 4.30 (m, 1H), 7.10-7.50 (m, 9H), 9.6 (br, 1H).

Compound 10

MS: m/z 322 (M+1).

¹H NMR (CDCl₃): d 1.60 (m, 4H), 1.90 (m, 2H), 2.60 (m, 2H), 3.10 (m, 1H), 3.84 (s, 2H), 4.50 (m, 1H), 7.10-7.50 (m, 9H), 9.6 (br, 1H).

2 g of Pd(OH)₂ was added into a solution of 30 ml of methanol containing 4.7 g of compound 10. The resulting suspension was hydrogenated at 50 psi for 12 hrs at room temperature. TLC indicated the reaction was complete over night. The solution was filtered through a pad of celite to remove the catalyst. The celite was washed with methanol twice (20 ml). The organics were combined and solvent was removed to give a pale solid which was purified by chromatography (10% MeOH, 90% EtOAc) to give an off white product 11 (1.79 g, 50.7%).

Compound 11

MS: m/z 232 (M+1).

¹H NMR (CDCl₃): d 1.50-1.85 (m, 8H), 2.60 (m, 2H), 4.30 (m, 1H), 7.10 (m, 3H), 7.30 (m, 1H).

To a mixture of 11 (1.7 g, 7.4 mmole), acetaldehyde (0.33 g, 7.4 mmole) in 50 ml of dichloroethane, 2.2 g of sodium triacetoxyborohydride (10.36 mmol, 1.4 eq) was added in one portion. Gas evolves between 30 min and 1 hr. The mixture was stirred over night. TLC indicated the reaction was complete. 1 N NaOH (500 ml) was added to quench the reaction. The organic layer was separated and the aqueous layer was extracted with EtOAc (300 ml x2). The combined organics were dried over potassium carbonate and the solvent was evaporated to give a brown oil which was chromatographed (10%Et₃N, 40%EtOAc and 50%Hexane) to give 1.5 g of product 2 as a sticky oil which recrystallized from TBME to give a white solid (78%).

Compound 2

MS: m/z 259.7 (M+1).


¹H NMR (CDCl₃): δ 1.15 (t, 3H), 1.50–1.95 (m, 6H), 2.40–2.75 (m, 4H), 2.95 (m, 1H), 4.35 (m, 1H), 7.10 (m, 3H), 7.35 (m, 1H).

1.5 g of Pd(OH)₂ was added into a solution of 30 ml of methanol containing 3.01 g of compound 9. The resulting suspension was hydrogenated at 50 psi for 12 hrs at room temperature. TLC indicated the reaction was complete over night. The solution was filtered through a pad of celite to remove the catalyst. The celite was washed with methanol twice (20 ml). The organics were combined and solvent was removed to give a pale solid which was purified by chromatography (10% MeOH, 90% EtOAc) to give an off white product 1 (1.68 g, 77.4%).

Compound 1

MS: m/z 232 (M+1).

¹H NMR (CDCl₃): δ 1.50 (m, 2H), 1.90–2.35 (m, 6H), 3.00 (m, 1H), 4.30 (m, 1H), 7.10–7.30 (m, 4H).

SCHEME 2:**Procedure:**

About 2.5 g of NaH was washed by THF twice, suspended in 100 ml of DMF, then 8.15 g of 7 (38 mmole) was added to the mixture. Gas evolves, and after 5 minutes, 7.13 g of ethyl iodide (45.7 mmole) was added. The mixture was stirred over night. LC/MS indicated that the starting material was completely consumed. The reaction was cooled down and H_2O was added to the mixture. The product started to precipitated out of solution. The crystals was collected by filtration to give 9.7 g of 12 (84.7%).

Compound 12

MS: m/z 303.3 ($\text{M}+1$).

^1H NMR (CDCl_3): d 1.30 (t, 3H), 1.70 – 1.90 (m, 6H), 2.50 (m, 2H), 3.85-4.00 (m, 6H), 4.50 (m, 1H), 7.05 (m, 3H), 7.25 (m, 1H).

A mixture of 12 (9.7 g, 32.2 mmole) and 9.72 g of PPTS in 50 ml of acetone and H_2O

(10:1) was refluxed over night. The mixture was cooled to room temperature and acetone was evaporated. Addition of water to the mixture initiated crystallization to give 6.85 g of product 13 (82.3%).

Compound 13

MS: m/z 259 ($M+1$).

^1H NMR (CDCl_3): δ 1.35 (t, 3H), 2.20 (m, 2H), 2.60 (m, 6H), 3.95 (q, 2H), 4.85 (m, 1H), 7.10 (m, 4H).

To a mixture of 13 (6.85 g, 26.5 mmole), benzylamine (2.84 g, 26.5 mmole), acetic acid (1.59 g, 26.5 mmole) in 150 ml of dichloroethane, 7.86 g of sodium triacetoxyborohydride (37.1 mmol, 1.4 eq) was added in one portion. Gas evolves between 30 min and 1 hr. The mixture was stirred over night. TLC indicated the reaction was complete. 1 N NaOH (500 ml) was added to quench the reaction. The organic layer was separated and the aqueous layer was extracted with EtOAc (300 ml x2). The combined organics were dried over potassium carbonate and the solvent was evaporated to give a brown solid, which was column filtrated (5%Et₃N, 25%EtOAc and 70%Hexane to 10%Et₃N, 40%EtOAc and 50%Hexane) to give 1.52g of product 14 as a white solid and 1.08 g of product 15 as a white solid.

Compound 14

MS: m/z 350 ($M+1$).

^1H NMR (CDCl_3): δ 1.35 (t, 3H), 1.50 (m, 2H), 1.65 (m, 4H), 1.95 (m, 2H), 2.60 (m, 2H), 3.02 (m, 1H), 3.83 (s, 2H), 3.95 (ddd, 2H), 4.45 (m, 1H), 7.00–7.50 (m, 9H).

Compound 15

MS: m/z 350 ($M+1$).

^1H NMR (CDCl_3): δ 1.35 (m, 5H), 1.90 (m, 2H), 2.10–2.35 (m, 4H), 2.70 (m, 1H), 3.83 (s, 2H), 3.95 (ddd, 2H), 4.40 (m, 1H), 7.00–7.50 (m, 9H).

0.3 g of Pd(OH)₂ was added into a solution of 20 ml of methanol containing 0.5 g of compound 14. The resulting suspension was hydrogenated at 50 psi for 12 hr at room temperature. TLC indicated the reaction was complete over night. The solution was filtered through a pad of celite to remove the catalyst. The celite was washed with methanol twice (20 ml). The organics were combined and solvent was removed to give a pale solid which was

purified by chromatography (10% MeOH, 90% EtOAc) to give an off white product 3 (300 mg, 50%).

Compound 3


MS: m/z 232 (M+1).

^1H NMR (CDCl_3): d 1.35 (t, 3H), 1.50–1.85 (m, 8H); 2.60 (m, 2H), 3.20 (m, 1H), 3.95 (ddd, 2H), 4.30 (m, 1H), 7.10 (m, 3H), 7.30 (m, 1H).

EXAMPLE 11

ATTACHMENT OF TAIL GROUPS

Tail groups were attached to the head groups according to the following procedures:



General procedure for alkylation:

To a solution of the amine (1 eq) and triethylamine (1 eq) in dimethylformamide, was added 1 eq of alkyl bromide or chloride in one portion. The mixture was stirred and heated at 80°C over night. TLC indicated the reaction was complete. The reaction was quenched by the addition of water followed by 1 N NaOH to pH 10. The mixture was extracted 2x with Et_2O . The combined organic extracts were dried over potassium carbonate and the solvent evaporated, followed by chromatography to give the pure product.

General procedure for reductive amination:

To a mixture of ketone or aldehyde (1 eq), amine (1 eq), and acetic acid (1 eq) in methanol, was added sodium cyanoborohydride (1.4 eq) in one portion. The mixture was stirred over night at room temperature. TLC indicated the reaction was complete. The reaction was quenched by the addition of water followed by 1 N NaOH to pH 10. The mixture was extracted 2x with Et₂O. The combined organic extracts were dried over potassium carbonate and the solvent evaporated, followed by chromatography to give the pure product.

The following compounds were prepared by attaching the tail groups using the general procedures described:

1-[4-(benzylamino)-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one

1-[4-(benzylamino)-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one

1-[4-[(naphth-2-yl-methyl)ethylamino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one

MS: m/z 400.2 (M+1)

1-[4-(norbornan-2-ylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one

MS: m/z 326.3 (M+1)

1-[4-[[4-(1-methylethyl)-cyclohexyl]amino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one

MS: m/z 356.4 (M+1)

1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one

MS: m/z 368.2 (M+1)

1-[4-(ethylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one

1-[4-(benzylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one

1-[4-(benzylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one

1-[4-[(indan-2-yl)benzylamino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one

MS: m/z 466.3 (M+1)

¹H-NMR (CDCl₃): d 1.30 (t, 3H), 1.50-1.75 (m, 2H), 1.90 (b, 2H), 2.02 (b, 2H), 2.20 (m, 2H), 2.80 (m, 1H), 2.99 (m, 4H), 3.75 (s, 2H), 3.90 (m, 3H), 4.25 (m, 1H), 6.95-7.45 (m, 13H).

1-[4-[(cyclooctylmethyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 99%

MS: m/z 384.5

¹H NMR (CDCl₃): d 1.40 -1.90 (m, 24H), 2.30 (m, 2H), 2.50(m, 2H), 2.90(m, 1H), 3.90(ddd, 2H), 4.20(m, 1H), 7.10(m, 3H), 7.30(m, 1H).

1-[4-[(naphth-2-yl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 97%

MS: m/z 399

¹H NMR (CDCl₃): d 1.50 (t, 3H), 1.80 (m, 5H), 2.0 (m, 2H), 2.70(m, 2H), 3.10(m, 1H), 3.90(m, 2H), 4.0 (m, 2H), 4.40(m, 1H), 7.10(m, 3H), 7.50(m, 4H), 7.90(m, 4H).

1-[4-[(*p*-benzyloxybenzyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 97%

MS: m/z 455

¹H NMR (CDCl₃): d 1.40 (t, 3H), 1.70 (m, 2H), 1.90 (m, 3H), 2.60(m, 4H), 3.10(m, 1H), 3.80(s, 2H), 4.0(m, 2H), 4.50(m, 1H), 5.10(s, 2H), 7.10(m, 6H), 7.50(m, 6H), 7.90(m, 1H).

1-[4-[(cyclooctylmethyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 99%

MS: m/z 369

¹H NMR (CDCl₃): d 1.40 (t, 3H), 1.70(m, 5H), 1.90(m, 12H), 2.10(m, 3H), 2.40(m, 2H), 2.50(d, 2H), 3.30(m, 1H), 3.90(m, 2H), 4.20(m, 1H), 7.10(m, 1H), 7.30(m, 3H).

1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one
LC: 99%

MS: m/z 395

¹H-NMR (CDCl₃): δ 1.40 (t, 3H), 1.70(m, 3H), 1.80(m, 3H), 1.90(m, 12H), 2.20(m, 2H), 2.30(m, 3H), 2.50(q, 2H), 3.10(m, 1H), 3.90(m, 2H), 4.20(m, 1H), 4.30(m, 1H), 7.0(m, 1H), 7.30(m, 3H).

1-[4-[(*p*-phenylbenzyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one
LC: 100%

MS: m/z 440.8 (M+1)

¹H-NMR (MeOH-d₄): δ 1.75 (m, 2H), 2.00 (m, 2H), 2.40-2.55 (m, 4H), 3.35-3.52 (m, 2H), 4.35 (s, 2H), 7.40 (m, 2H), 7.59 (t, 2H), 7.60-7.72 (m, 6H), 7.78 (d, 2H).

1-[4-[(1,2,3,4-tetrahydronaphthyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 93.9%

MS: m/z 405.7 (M+1)

¹H-NMR (MeOH-d₄): δ 1.70 (m, 2H), 1.85 (m, 1H), 2.02 (m, 2H), 2.39 (b, 3H), 2.50 (m, 2H), 2.90 (m, 1H), 3.00 (b, 2H), 3.35 (m, 1H), 3.60 (m, 1H), 3.72 (b, 1H), 4.35 (m, 1H), 7.15 (b, 4H), 7.40 (d, 1H), 7.60 (s, 1H), 7.65 (d, 1H).

1-[4-[(4-propyl-cyclohexyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 100%

MS: m/z 399.6 (M+1)

¹H-NMR (MeOH-d₄): δ 0.95 (t, 3H), 1.10 (m, 1H), 1.20-1.60 (m, 6H), 1.70 (b, 5H), 1.80-2.00 (m, 4H), 2.10 (m, 1H), 2.30 (b, 2H), 2.45 (m, 2H), 3.25 (m, 1H), 3.50 (m, 1H), 4.40 (m, 1H), 7.40 (d, 1H), 7.60 (s, 1H), 7.65 (d, 1H).

1-[4-[(5-methylhex-2-yl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one
LC: 100%

MS: m/z 373.5 (M+1)

¹H-NMR (MeOH-d₄): d 0.95 (d, 6H), 1.25-1.40 (m, 5H), 1.50-1.75 (m, 4H), 1.85 (m, 1H), 1.95 (b, 2H), 2.30 (m, 2H), 2.40-2.55 (m, 2H), 3.35-3.55 (m, 2H), 4.38 (m, 1H), 7.40 (d, 1H), 7.60 (s, 1H), 7.70 (d, 1H).

1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 100%

MS: m/z 411.7 (M+1)

¹H-NMR (MeOH-d₄): d 0.90-2.10 (m, 18H), 2.10-2.50 (m, 5H), 2.82 (m, 1H), 3.50 (m, 2H), 4.35 (m, 1H), 7.42 (d, 1H), 7.60 (s, 1H), 7.70 (d, 1H).

1-[4-(cyclooctylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;

LC: 95.4%

MS: m/z 385.7 (M+1)

¹H-NMR (MeOH-d₄): d 1.50-2.10 (m, 13H), 2.30 (m, 2H), 2.40-2.52 (m, 3H), 2.80-2.95 (m, 3H), 3.45 (m, 2H), 3.70 (m, 1H), 4.38 (m, 1H), 7.40 (d, 1H), 7.63 (s, 1H), 7.70 (d, 1H).

1-[4-[(indan-2-yl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 100%

MS: m/z 391.6 (M+1)

¹H-NMR (MeOH-d₄): d 1.70 (m, 2H), 2.00 (m, 2H), 2.40-2.60 (m, 4H), 3.10-3.20 (m, 2H), 3.50 (m, 3H), 4.30-4.45 (m, 2H), 7.25 (m, 2H), 7.35 (m, 2H), 7.42 (d, 1H), 7.60 (s, 1H), 7.72 (d, 1H).

1-[4-(benzylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 100%

MS: m/z 399.5 (M+1)

¹H-NMR (MeOH-d₄): d 1.40-1.85 (m, 15H), 2.00 (m, 4H), 2.25-2.50 (m, 4H), 2.93 (d, 2H), 3.30 (m, 1H), 4.30 (m, 1H), 7.36 (d, 1H), 7.60 (s, 1H), 7.65 (d, 1H).

1-[4-[(4-phenyl-cyclohexyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 100%

MS: m/z 433.7 (M+1)

¹H-NMR (MeOH-d₄): δ 1.65 (m, 2H), 1.85-2.20 (m, 8H), 2.25-2.50 (m, 5H), 3.90 (m, 1H), 3.50 (m, 2H), 3.58 (m, 1H), 4.30 (m, 1H), 7.15-7.40 (m, 6H), 7.60 (s, 1H), 7.65 (d, 1H).

1-[4-(dibenzylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 100%

MS: m/z 455.6 (M+1)

¹H-NMR (MeOH-d₄): δ 2.00-2.25 (m, 4H), 2.40 (m, 4H), 3.52 (m, 2H), 4.25-4.65 (m, 4H), 7.30 (d, 1H), 7.45-7.58 (m, 10H), 7.60 (s, 1H), 7.65 (d, 1H).

1-[4-[(5-methylhex-2-yl)amino]-cyclohexyl]-7-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one

LC: 99.1%

MS: m/z 373.3 (M+1)

¹H-NMR (MeOH-d₄): δ 0.95 (d, 6H), 1.30 (d, 3H), 1.45-1.68 (m, 5H), 1.75 (m, 1H), 2.00 (m, 2H), 2.18-2.32 (m, 3H), 2.60 (m, 2H), 3.20-3.40 (m, 2H), 4.30 (m, 1H), 7.05-7.20 (m, 3H).

Other compounds within the scope of formula (III) or (IIIA) of the present invention can be synthesized by analogous techniques.

EXAMPLE 12

Nociceptin affinity at the ORL1 receptor for preferred compounds was obtained using the following assay:

Membranes from recombinant HEK-293 cells expressing the human opioid receptor-like receptor (ORL-1) (Receptor Biology) were prepared by lysing cells in ice-cold hypotonic buffer (2.5 mM MgCl₂, 50 mM HEPES, pH 7.4) (10 ml/10 cm dish) followed by homogenization with a tissue grinder/teflon pestle. Membranes were collected by centrifugation at 30,000 x g for 15 min at 4°C and pellets resuspended in hypotonic buffer to a final concentration of 1-3 mg/ml. Protein concentrations were determined using the BioRad protein assay reagent with bovine

serum albumen as standard. Aliquots of the ORL-1 receptor membranes were stored at -80°C .

Functional SGTPgS binding assays were conducted as follows. ORL-1 membrane solution was prepared by sequentially adding final concentrations of 0.066 mg/ml ORL-1 membrane protein, 10 mg/ml saponin, 3 mM GDP and 0.20 nM [^{35}S]GTPgS to binding buffer (100 mM NaCl, 10 mM MgCl_2 , 20 mM HEPES, pH 7.4) on ice. The prepared membrane solution (190 μl /well) was transferred to 96-shallow well polypropylene plates containing 10 μl of 20x concentrated stock solutions of agonist prepared in DMSO. Plates were incubated for 30 min at room temperature with shaking. Reactions were terminated by rapid filtration onto 96-well Unifilter GF/B filter plates (Packard) using a 96-well tissue harvester (Brandel) and followed by three filtration washes with 200 μl ice-cold binding buffer (10 mM NaH_2PO_4 , 10 mM Na_2HPO_4 , pH 7.4). Filter plates were subsequently dried at 50°C for 2-3 hours. Fifty μl /well scintillation cocktail (BetaScint; Wallac) was added and plates were counted in a Packard Top-Count for 1 min/well.

Data was analyzed using the curve fitting functions in GraphPad PRISM[®], v. 3.0 and the results are set forth in table 3 below:


TABLE 3	
Nociceptin Affinity	
Compound	calc K_i (nM)
3-ethyl-1-(<i>p</i> -phenylbenzyl)-1,3-dihydro-2H-benzimidazol-2-one	509
3-ethyl-1-(5-methylhex-2-yl)-1,3-dihydro-2H-benzimidazol-2-one	23
3-ethyl-1-(4-propylcyclohexyl)-1,3-dihydro-2H-benzimidazol-2-one	68
3-ethyl-1-(decahydro-2-naphthyl)-1,3-dihydro-2H-benzimidazol-2-one	1.6
3-ethyl-1-(naphth-2-yl-methyl)-1,3-dihydro-2H-benzimidazol-	198

2-one	
1-(<i>p</i> -benzyloxybenzyl)-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	438
1-benzyl-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	296
1-[4-(benzylamino)-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	trans: 112 cis: >10,000
3-ethyl-1-(naphthylmethyl)-1,3-dihydro-2H-benzimidazol-2-one	39
3-ethyl-1-[5-(3-fluorophenyl)-5-(4-fluorophenyl)-hexyl]-1,3-dihydro-2H-benzimidazol-2-one	148
1-[4-[(naphth-2-yl-methyl)ethylamino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one	3598
1-[4-(norbornan-2-ylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[[4-(1-methylethyl)-cyclohexyl]amino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-(ethylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one	9179
1-[4-(benzylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one	trans: 273 cis: >10000
1-[4-[(indan-2-yl)benzylamino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[(cyclooctylmethyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	115
1-[4-[(naphth-2-yl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	961
1-[4-[(<i>p</i> -benzyloxybenzyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	2935
1-[4-[(cyclooctylmethyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	286
1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	288
1-[4-(benzylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-(dibenzylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[(<i>p</i> -phenylbenzyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[(1,2,3,4-tetrahydronaphthyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[(4-propyl-cyclohexyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000

1-[4-[(5-methylhex-2-yl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-(cyclooctylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[(indan-2-yl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[(4-phenyl-cyclohexyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000
1-[4-[(5-methylhex-2-yl)amino]-cyclohexyl]-7-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one	>10000

EXAMPLE 13

SYNTHESIS OF SUBSTITUTED BENZIMIDAZOLE HEAD GROUPS.

Procedure:

Sodium hydride 60% dispersion in mineral oil (0.67 g, 16.7 mmol) was washed with dry pentane and then suspended in 80 mL of dry THF under N₂. Compound 1 (European patent 0029707) (3.80 g, 11.1 mmol) was added, the mixture stirred at room temperature for 15 min and then warmed to 50°C. Ethyl bromide (1.06 mL, 13.3 mmol) was added and the resulting mixture stirred at 50°C for 18 hr. TLC (SiO₂, CH₂Cl₂:MeOH 96:4) showed that the reaction was *ca* 40% complete. Additional sodium hydride (0.67 g) and ethyl bromide (1.06 mL) were added. After heating at 50°C for an additional 24 hr the reaction mixture was cooled to room temperature and quenched with water. The layers were separated and the aqueous layer extracted with ethyl

acetate (1x). The combined organic extracts were washed with aqueous sodium bicarbonate solution (1x), dried over MgSO_4 and the solvent was evaporated to give the crude product as a yellow solid. Trituration with diethyl ether gave pure 2 as a white solid (3.38 g, 82%).

$^1\text{H-NMR}$ (CDCl_3): d 1.45-1.55 (m, 12H), 1.82 (bs, 2H), 2.30 (m, 2H), 2.87 (m, 2H), 4.30 (bs, 2H), 4.41 (q, 2H), 4.82 (m, 1H), 7.10-7.30 (m, 4H).


To a solution of 2 (3.60 g, 9.74 mmol) in 100 mL of ethyl acetate was added a 25 mL of a 1:1 mixture of ethyl acetate and concentrated HCl . The mixture was stirred vigorously at room temperature for 2hr. and evaporated to dryness. The residue was neutralized with 50 mL of methanolic ammonia 10:1 and again evaporated to dryness. The residue was suspended in 100 mL a 1:1 mixture of MeOH and CH_2Cl_2 , filtered and the filtrate evaporated to dryness to leave an off-white solid. Flash chromatography on silica gel, eluting with $\text{CH}_2\text{Cl}_2:\text{MeOH}:\text{NH}_3$ (300:10:1) gave pure 3 as a white crystalline solid (1.98 g, 76%).

$^1\text{H-NMR}$ (CDCl_3): d 1.45 (t, 3H), 1.82 (bs, 2H), 2.33 (m, 2H), 2.80 (m, 2H), 4.40 (q, 2H), 4.80 (m, 1H), 7.10-7.30 (m, 3H), 7.45 (d, 1H).

EXAMPLE 14

ATTACHMENT OF TAIL GROUPS

Tail groups were attached to the head groups according to the following procedures:



General procedure for alkylation:

To a solution of the amine (1 eq) and triethylamine (1 eq) in dimethylformamide, was added 1 eq of alkyl bromide or chloride in one portion. The mixture was stirred and heated at 80°C over night. TLC indicated the reaction was complete. The reaction was quenched by the addition of water followed by 1 N NaOH to pH 10. The mixture was extracted 2x with Et₂O. The combined organic extracts were dried over potassium carbonate and the solvent evaporated, followed by chromatography to give the pure product.

General procedure for reductive amination:

To a mixture of ketone or aldehyde (1 eq), amine (1 eq), and acetic acid (1 eq) in methanol, was added sodium cyanoborohydride (1.4 eq) in one portion. The mixture was stirred over night at room temperature. TLC indicated the reaction was complete. The reaction was quenched by the addition of water followed by 1 N NaOH to pH 10. The mixture was extracted 2x with Et₂O. The combined organic extracts were dried over potassium carbonate and the solvent evaporated, followed by chromatography to give the pure product.

The following compounds were prepared by attaching the tail groups using the general procedures described:

2-cyanoimino-3-ethyl-1-[1-(*p*-phenylbenzyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole
¹H-NMR (CDCl₃): d 1.50 (t, 3H), 1.88 (m, 2H), 2.28 (m, 2H), 2.62 (m, 2H), 3.12 (m, 2H), 3.65 (s, 2H), 4.48 (q, 2H), 4.80 (m, 1H), 7.15-7.70 (m, 13H).

2-cyanoimino-3-ethyl-1-[1-(*p*-benzyloxybenzyl)-4-piperidinyl] 1,3-dihydro-2H-benzimidazole

LC: 96.5%

MS: m/z 466.5 (M+1)

¹H-NMR (CDCl₃): d 1.55 (t, 3H), 1.82 (m, 2H), 2.25 (m, 2H), 2.50 (m, 2H), 3.10 (m, 2H), 3.55 (s, 2H), 4.48 (q, 2H), 4.78 (m, 1H), 5.20 (s, 2H), 7.00 (d, 2H), 7.15-7.65 (m, 11H).

2-cyanoimino-3-ethyl-1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 93.9%

MS: m/z

¹H-NMR (CDCl₃): d 1.55 (t, 3H), 1.80 (m, 2H), 2.30 (t, 2H), 2.52 (m, 2H), 3.18 (bd, 2H), 3.78 (s, 2H), 4.50 (q, 2H), 4.80 (m, 1H), 7.20-7.90 (m, 11H).

2-cyanoimino-3-ethyl-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

MS: m/z 394.4 (M+1)

¹H-NMR (CDCl₃): d 0.90-2.28 (m, 21H), 3.10 (m, 4H), 3.62 (m, 2H), 4.42 (q, 2H), 5.15 (m, 1H), 7.20 (d, 1H), 7.30 (m, 1H), 7.50 (t, 1H), 7.80 (b, 1H).

2-cyanoimino-3-ethyl-1-[1-[4-(2-propyl)-cyclohexyl]-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 100%

MS: m/z 394.5 (M+1)

¹H-NMR (CDCl₃): d 0.90 (d, 3H), 0.98 (d, 3H), 1.15-2.35 (m, 14H), 3.10 (m, 5H), 3.70 (m, 2H), 3.92 (bs, 1H), 4.40 (q, 2H), 5.20 (m, 1H), 7.20 (d, 1H), 7.38 (d, 1H), 7.52 (t, 1H), 7.80 (m, 1H).

2-cyanoimino-3-ethyl-1-[1-(decahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 93.9%

MS: m/z 406.6 (M+1)

¹H-NMR (CDCl₃): d 1.25-2.35 (m, 24H), 1.15 (m, 4H), 3.60 (m, 2H), 4.40 (m, 2H), 4.20 (m, 1H), 7.20-7.80 (m, 4H).

2-cyanoimino-3-ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 100%

MS: m/z 380.3 (M+1)

¹H-NMR (CDCl₃): d 1.50-1.80 (m, 13H), 1.90 (m, 2H), 2.10 (m, 4H), 3.05 (m, 3H), 3.30 (m, 1H), 3.45 (m, 2H), 3.90 (m, 1H), 4.42 (q, 2H), 5.15 (m, 1H), 7.20 (d, 1H), 7.35 (d, 1H), 7.50 (m, 1H), 7.78 (m, 1H).

2-cyanoimino-3-ethyl-1-[1-(10,11-dihydro-5H-dibenzo[a,d]-cyclohepten-5-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 94.5%

MS: m/z 462.2 (M+1)

¹H-NMR (CDCl₃): d 1.40 (t, 3H), 1.70 (bs, 2H), 2.01 (m, 2H), 2.28 (m, 2H), 2.80 (m, 4H), 3.95 (s, 1H), 4.02 (m, 2H), 4.32 (q, 2H), 4.65 (m, 1H), 7.00-7.32 (m, 12H).

2-cyanoimino-3-ethyl-1-[1-(3,3-Bis(phenyl)propyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

MS: m/z 464.2 (M+1)

¹H-NMR (CDCl₃): d 1.40 (t, 3H), 1.73 (bs, 2H), 2.09 (m, 2H), 2.18-2.45 (m, 6H), 2.98 (b, 2H), 3.93 (t, 1H), 4.35 (q, 2H), 4.63 (m, 1H), 7.10-7.30 (m, 13H), 7.40 (d, 1H).

2-cyanoimino-3-ethyl-1-[1-(1,2,3,4-tetrahydronaphthyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 94.0%

MS: m/z 400.2 (M+1)

¹H-NMR (CDCl₃): d 1.30-1.70 (m, 6H), 1.85 (m, 2H), 2.05 (m, 1H), 2.45 (m, 3H), 2.85 (m, 4H), 3.10 (m, 2H), 4.35 (q, 2H), 4.71 (m, 1H), 7.00-7.60 (m, 8H).

2-cyanoimino-3-ethyl-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 94.9%

MS: m/z 368.3 (M+1)

¹H-NMR (CDCl₃): d 0.85 (d, 6H), 0.95 (d, 3H), 1.12-1.65 (m, 8H), 1.80 (m, 2H), 2.27-2.60 (m, 5H), 2.85 (m, 2H), 4.38 (m, 2H), 4.62 (m, 1H), 7.08-7.30 (m, 3H), 7.45 (m, 1H).

2-cyanoimino-3-ethyl-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 99.2%

MS: m/z 364.7 (M+1)

¹H-NMR (CDCl₃): d 1.10-2.10 (m, 13H), 2.35 (m, 1H), 2.50-2.70 (m, 3H), 2.70-2.90 (m, 3H), 3.50 (m, 2H), 4.50 (q, 2H), 4.80 (m, 1H), 7.35 (m, 2H), 7.48 (m, 1H), 7.75 (m, 1H).

2-cyanoimino-3-ethyl-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 92.1 %

MS: m/z 386.2 (M+1)

¹H-NMR (CDCl₃): d 1.42 (t, 3H), 1.82 (m, 2H), 2.21 (m, 2H), 2.43 (m, 2H), 2.88 (m, 2H), 3.02-3.19 (m, 4H), 3.23 (m, 1H), 4.38 (q, 2H), 4.80 (m, 1H), 7.08-7.30 (m, 7H), 7.45 (d, 1H).

2-cyanoimino-3-ethyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 100%

MS: m/z 394.7 (M+1)

¹H-NMR (MeOH): d 1.35-2.00 (m, 20H), 2.60-2.85 (m, 6H), 3.40 (m, 2H), 2.52 (q, 2H), 4.90 (m, 1H), 7.35 (m, 2H), 7.48 (m, 1H), 7.70 (m, 1H).

2-cyanoimino-3-(2-hydroxy)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 100%

MS: m/z 396.3 (M+1)

¹H-NMR (DMSO): 7.52 (dt, 1H), 7.45 (dt, 1H), 7.21 (m, 2H), 4.97 (t, 1H), 4.55 (m, 1H), 4.38 (t, 2H), 3.76 (q, 2H), 2.88 (m, 2H), 2.61 (bt, 1H), 2.33 (m, 4H), 1.76-1.37 (m, 16H).

2-cyanoimino-3-methoxycarbonylmethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 98.3%

MS: m/z 424.2 (M+1)

¹H-NMR (DMSO): 7.56 (dd, 1H), 7.51 (dd, 1H), 7.25 (m, 2H), 5.26 (s, 2H), 4.56 (m, 1H), 3.72 (s, 3H), 3.34 (m, 2H), 2.78 (m, 2H), 2.62 (bt, 1H), 2.32 (m, 4H), 1.80-1.35 (m, 16H).

2-cyanoimino-3-cyanomethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole
LC: 100%

MS: m/z 391.2 (M+1)

¹H-NMR (DMSO): 7.60 (m, 2H), 7.31 (m, 2H), 5.48 (s, 2H), 4.77 (m, 1H), 3.33 (d, 2H), 2.88 (m, 2H), 2.62 (bt, 1H), 2.33 (m, 4H), 1.86-1.37 (m, 16H).

2-cyanoimino-3-butyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole
LC: 95.4%

MS: m/z 352.2 (M+1)

¹H-NMR (DMSO): 7.58 (dd, 1H), 7.49 (dd, 1H), 7.24 (m, 2H), 6.55 (s, 2H), 4.59 (m, 1H), 4.34 (t, 2H), 2.97 (m, 2H), 2.80 (m, 1H), 2.55 (m, 2H), 2.38 (m, 2H), 1.80-1.30 (m, 18H), 0.90 (t, 3H).

2-cyanoimino-3-(2-methanesulfonamido)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 100%

MS: m/z 473.2 (M+1)

¹H-NMR (DMSO): 7.53 (dd, 1H), 7.44 (dd, 1H), 7.23 (m, 2H), 4.60 (m, 1H), 4.35 (t, 2H), 3.37 (t, 2H), 2.87 (m, 2H), 2.82 (s, 3H), 2.60 (bt, 1H), 2.31 (m, 4H), 1.76-1.37 (m, 15H).

2-cyanoimino-3-acetomido-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole
LC: 100%

MS: m/z 409.2 (M+1)

¹H-NMR (DMSO): 7.75 (s, 1H), 7.52 (dd, 1H), 7.37 (s, 1H), 7.30 (dd, 1H), 7.20 (m, 2H), 4.96 (s, 2H), 4.55 (m, 1H), 3.33 (d, 2H), 2.88 (m, 2H), 2.62 (bt, 1H), 2.30 (m, 4H), 1.80-1.37 (m, 15H).

2-cyanoimino-3-carboxymethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole
LC: 97.5%

MS: m/z 409.9 (M+1)

¹H-NMR (DMSO): 7.45 (dd, 1H), 7.14 (m, 3H), 4.57 (s, 2H), 4.50 (m, 1H), 2.87 (m, 2H), 2.61 (bt, 1H), 2.33 (m, 4H), 1.75-1.37 (m, 15H).

2-cyanoimino-3-(2-dimethylamino)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

LC: 100%

MS: m/z 423.3 (M+1)

¹H-NMR (DMSO): 7.60-6.96 (m, 4H), 6.54 (2H, s), 4.65 (m, 1H), 4.40 (t, 2H), 3.90 (t, 2H), 3.05 (m, 4H), 2.90 (m, 1H), 2.63 (m, 3H), 2.56-2.37 (m, 4H), 1.85-1.35 (m, 15H).

2-cyanoimino-1-[1-(cyclooctyl)-3-hydroxymethyl-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

2-cyanoimino-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-7-azabenzimidazole;

2-cyanoimino-1-[1-(cyclooctyl)-2,6-ethano-4-one-4-piperidinyl]-1,3-dihydro-2H-benzimidazole

Other compounds within the scope of formula (IV) or (IVA) of the present invention can be synthesized by analogous techniques.

EXAMPLE 15

Nociceptin affinity at the ORL1 receptor for preferred compounds was obtained using the following assay:

Membranes from recombinant HEK-293 cells expressing the human opioid receptor-like receptor (ORL-1) (Receptor Biology) were prepared by lysing cells in ice-cold hypotonic buffer (2.5 mM MgCl₂, 50 mM HEPES, pH 7.4) (10 ml/10 cm dish) followed by homogenization with a tissue grinder/teflon pestle. Membranes were collected by centrifugation at 30,000 x g for 15 min at 4°C and pellets resuspended in hypotonic buffer to a final concentration of 1-3 mg/ml. Protein concentrations were determined using the BioRad protein assay reagent with bovine serum albumen as standard. Aliquots of the ORL-1 receptor membranes were stored at -80°C.

Functional SGTPgS binding assays were conducted as follows. ORL-1 membrane solution was prepared by sequentially adding final concentrations of 0.066 mg/ml ORL-1 membrane protein, 10 mg/ml saponin, 3 mM GDP and 0.20 nM [³⁵S]GTPgS to binding buffer (100 mM NaCl, 10 mM MgCl₂, 20 mM HEPES, pH 7.4) on ice. The prepared membrane solution (190 ml/well) was transferred to 96-shallow well polypropylene plates containing 10 ml of 20x concentrated stock solutions of agonist prepared in DMSO. Plates were incubated for 30 min at room temperature with shaking. Reactions were terminated by rapid filtration onto 96-well Unifilter GF/B filter plates (Packard) using a 96-well tissue harvester (Brandel) and followed by three filtration washes with 200 ml ice-cold binding buffer (10 mM NaH₂PO₄, 10 mM Na₂HPO₄, pH 7.4). Filter plates were subsequently dried at 50°C for 2-3 hours. Fifty ml/well scintillation cocktail (BetaScint; Wallac) was added and plates were counted in a Packard Top-Count for 1 min/well.

Data was analyzed using the curve fitting functions in GraphPad PRISM[®], v. 3.0 and the results are set forth in table 4 below:

TABLE 45	
Nociceptin Affinity	
Compound	calc K_i (nM)
2-cyanoimino-3-ethyl-1-[1-(<i>p</i> -phenylbenzyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	5558
2-cyanoimino-3-ethyl-1-[1-(<i>p</i> -benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	1660
2-cyanoimino-3-ethyl-1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	882
2-cyanoimino-3-ethyl-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	241
2-cyanoimino-3-ethyl-1-[1-[4-(2-propyl)-cyclohexyl]-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	6.9
2-cyanoimino-3-ethyl-1-[1-(decahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	6.6
2-cyanoimino-3-ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	5.57
2-cyanoimino-3-ethyl-1-[1-(10,11-Dihydro-5H-dibenzo[a,d]-cyclohepten-5-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;	> 10,000
2-cyanoimino-3-ethyl-1-[1-(3,3-Bis(phenyl)propyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;	80

2-cyanoimino-3-ethyl-1-[1-(1,2,3,4-tetrahydronaphthyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;	157
2-cyanoimino-3-ethyl-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;	76
2-cyanoimino-3-ethyl-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;	323
2-cyanoimino-3-ethyl-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole; and	89
2-cyanoimino-3-ethyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole.	7.1
2-cyanoimino-3-(2-hydroxy)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	6.4
2-cyanoimino-3-methoxycarbonylmethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	3.3
2-cyanoimino-3-cyanomethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	.97
2-cyanoimino-3-butyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	1.36
2-cyanoimino-3-(2-methanesulfonamido)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	78
2-cyanoimino-3-acetomido-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	11
2-cyanoimino-3-carboxymethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	201
2-cyanoimino-3-(2-dimethylamino)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	18
2-cyanoimino-1-[1-(cyclooctyl)-3-hydroxymethyl-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	473
2-cyanoimino-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-7-azabenzimidazole	3743
2-cyanoimino-1-[1-(cyclooctyl)-2,6-ethano-4-one-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	19

Example 16

Affinity at the μ receptor for compounds was obtained according to the following assay:

Mu opioid receptor membrane solution was prepared by sequentially adding final concentrations of 0.075 $\mu\text{g}/\mu\text{l}$ of the desired membrane protein, 10 $\mu\text{g}/\text{ml}$ saponin, 3 μM GDP and 0.20 nM [^{35}S]GTP γS to binding buffer (100 mM NaCl, 10 mM MgCl_2 , 20 mM HEPES, pH 7.4) on ice. The prepared membrane solution (190 $\mu\text{l}/\text{well}$) was transferred to 96-shallow well


polypropylene plates containing 10 μ l of 20x concentrated stock solutions of agonist prepared in DMSO. Plates were incubated for 30 min at room temperature with shaking. Reactions were terminated by rapid filtration onto 96-well Unifilter GF/B filter plates (Packard) using a 96-well tissue harvester (Brandel) and followed by three filtration washes with 200 μ l ice-cold binding buffer (10 mM NaH_2PO_4 , 10 mM Na_2HPO_4 , pH 7.4). Filter plates were subsequently dried at 50° C for 2-3 hours. Fifty μ l/well scintillation cocktail (MicroScint20, Packard) was added and plates were counted in a Packard Top-Count for 1 min/well.

Data were analyzed using the curve fitting functions in GraphPad PRISM™, v. 3.0 and the results for several compounds are set forth in table 5 below:

TABLE 5	
Mu Receptor Affinity	
Compound	calc K_i (nM)
3-[1-(naphth-1-yl-methyl)-4-piperidinyl]-2H-benzoxazol-2-one	340
3-[1-(3,3-diphenylpropyl)-4-piperidinyl]-2H-benzoxazol-2-one	726
3-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-2H-benzoxazol-2-one	343
3-[1-(4-propyl-cyclohexyl)-4-piperidinyl]-2H-benzoxazol-2-one	145
3-ethylidene-1-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	23.3
3-ethylidene-1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	137
3-ethylidene-1-[1-(<i>p</i> -benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	1150
3-ethylidene-1-[1-(3,3-diphenylpropyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one	24
1-[4-[(naphth-2-yl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one	2.1
2-cyanoimino-3-ethyl-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	46
2-cyanoimino-3-ethyl-1-[1-(1,2,3,4-tetrahydronaphthyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	458
2-cyanoimino-3-ethyl-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	15
2-cyanoimino-3-ethyl-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole	1653

What is claimed is:

1. A compound of the formula (I):



(I)

wherein

D is a 5-8 membered cycloalkyl, 5-8 membered heterocyclic or a 6 membered aromatic or heteroaromatic group;


n is an integer from 0 to 3;

A, B and Q are independently hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy, -CH₂OH, -NHSO₂, hydroxyC₁₋₁₀alkyl-, aminocarbonyl-, C₁₋₄alkylaminocarbonyl-, diC₁₋₄alkylaminocarbonyl-, acylamino-, acylaminoalkyl-, amide, sulfonylaminoC₁₋₁₀alkyl-, or A-B can together form a C₂₋₆ bridge, or B-Q can together form a C₃₋₇ bridge, or A-Q can together form a C₁₋₅ bridge;

Z is selected from the group consisting of a bond, straight or branched C₁₋₆ alkylene, -NH-, -CH₂O-, -CH₂NH-, -CH₂N(CH₃)-, -NHCH₂-, -CH₂CONH-, -NHCH₂CO-, -CH₂CO-, -COCH₂-, -CH₂COCH₂-, -CH(CH₃)-, -CH=, -O- and -HC=CH-, wherein the carbon and/or

nitrogen atoms are unsubstituted or substituted with one or more lower alkyl, hydroxy, halo or alkoxy group;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, $-COOV_1$, $-C_{1-4}COOV_1$, cyano, cyano C_{1-10} alkyl-, cyano C_{3-10} cycloalkyl-, NH_2SO_2 -, $NH_2SO_2C_{1-4}$ alkyl-, NH_2SOC_{1-4} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, di C_{1-4} alkylaminocarbonyl-, benzyl, C_{3-12} cycloalkenyl-, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a hetero-monocyclic ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ; and wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, or benzyl of R_1 is optionally substituted with 1-3 substituents selected from the group consisting of halogen, hydroxy, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, cyano, $-COOV_1$, $-C_{1-4}COOV_1$, cyano C_{1-10} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, a 5-membered heteroaromatic C_{0-4} alkyl-, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl-, C_{1-10} alkoxy-, and cyano; and wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, or spiro ring system of the formula (V) is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy or benzyloxy is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

W_1 is hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{1-10} alkoxy, C_{3-12} cycloalkoxy, $-CH_2OH$, amino, C_{1-4} alkylamino-, diC_{1-4} alkylamino-, or a 5-membered heteroaromatic ring optionally substituted with 1-3 lower alkyl;

V_1 is H, C_{1-6} alkyl, C_{3-6} cycloalkyl, benzyl or phenyl;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl- and halogen, said alkyl or cycloalkyl optionally substituted with an oxo, amino, alkylamino or dialkylamino group;

or a pharmaceutically acceptable salt thereof or a solvate thereof.


2. A compound of claim 1, wherein D is phenyl or a 6 membered heteroaromatic group containing 1-3 nitrogen atoms.

3. A compound of claim 1, wherein ZR_1 is cyclohexylethyl-, cyclohexylmethyl-, cyclopentylmethyl-, dimethylcyclohexylmethyl-, phenylethyl-, pyrrolyltrifluoroethyl-, thienyltrifluoroethyl-, pyridylethyl-, cyclopentyl-, cyclohexyl-, methoxycyclohexyl-, tetrahydropyranyl-, propylpiperidiny-, indolylmethyl-, pyrazoylpentyl-, thiazolyethyl-, phenyltrifluoroethyl-, hydroxyhexyl-, methoxyhexyl-, isopropoxybutyl-, hexyl-, or oxocanylpropyl-.

4. A compound of claim 1, wherein ZR_1 is $-CH_2COOV_1$, tetrazolylmethyl-, cyanomethyl-, NH_2SO_2 methyl-, NH_2SO methyl-, aminocarbonylmethyl-, C_{1-4} alkylaminocarbonylmethyl-, or diC_{1-4} alkylaminocarbonylmethyl-.

5. A compound of claim 1, wherein ZR_1 is 3,3 diphenylpropyl optionally substituted at the 3 carbon of the propyl with $-COOV_1$, tetrazolyl C_{0-4} alkyl-, cyano-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, or diC_{1-4} alkylaminocarbonyl-.

6. A compound of the formula (IA):




(IA)

wherein

n is an integer from 0 to 3;

Z is selected from the group consisting of a bond, -CH₂-, -NH-, -CH₂O-, -CH₂CH₂-, -CH₂NH-, -CH₂N(CH₃)-, -NHCH₂-, -CH₂CONH-, -NHCH₂CO-, -CH₂CO-, -COCH₂-, -CH₂COCH₂-, -CH(CH₃)-, -CH=, and -HC=CH-, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with a lower alkyl, halogen, hydroxy or alkoxy group;

R₁ is selected from the group consisting of hydrogen, C₁₋₁₀alkyl, C₃₋₁₂cycloalkyl, C₂₋₁₀alkenyl, amino, C₁₋₁₀alkylamino, C₃₋₁₂cycloalkylamino, benzyl, C₃₋₁₂cycloalkenyl, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a hetero-monocyclic ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ;

wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, or benzyl is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, cyano, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, and spiro ring system of the formula (V) are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy and benzyloxy are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

or a pharmaceutically acceptable salt thereof.


7. A compound of claim 6, wherein R_1 is alkyl selected from the group consisting of methyl, ethyl, propyl, butyl, pentyl and hexyl.
8. A compound of claim 6, wherein R_1 is cycloalkyl selected from the group consisting of cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, and norbornyl.
9. A compound of claim 6, wherein R_1 is tetrahydronaphthyl, decahydronaphthyl or dibenzocycloheptyl.
10. A compound of claim 6, wherein R_1 is phenyl or benzyl.
11. A compound of claim 6, wherein R_1 is a bicyclic aromatic ring.

12. A compound of claim 11, wherein said bicyclic aromatic ring is indenyl, quinoline or naphthyl.
13. A compound of claim 6, wherein Z is a bond, methyl, or ethyl.
14. A compound of claim 6, wherein n is 0.
15. A compound of claim 6, wherein X₁ and X₂ are both O.
16. A compound selected from the group consisting of:
 - 3-[1-(naphth-2-yl-methyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(naphth-1-yl-methyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(p-phenylbenzyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(p-benzyloxybenzyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(p-cyanobenzyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(3,3-diphenylpropyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-[4,4-Bis-(4-fluorophenyl)butyl]-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(2-phenylethyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(cyclooctylmethyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(5-methylhex-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(10,11-Dihydro-5H-dibenzo[a,d]-cyclohepten-5-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(4-propyl-cyclohexyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(norbornan-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(decahydro-2-naphthyl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(3,3-dimethyl-1,5-dioxaspiro[5.5]undeca-9-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-[4-(1-methylethyl)-cyclohexyl]-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-2H-benzoxazol-2-one;
 - 3-[1-(cyclooctyl)-4-piperidinyl]-2H-benzoxazol-2-one; and

pharmaceutically acceptable salts thereof.

17. A pharmaceutical composition comprising a compound of claim 1 and at least one pharmaceutically acceptable excipient.
18. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 1.
19. A method of modulating a pharmacological response from the ORL1 receptor comprising administering to a patient in need thereof an effective amount of a compound according to claim 1.
20. A pharmaceutical composition comprising a compound of claim 6 and at least one pharmaceutically acceptable excipient.
21. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 6.
22. A method of modulating a pharmacological response from the ORL1 receptor comprising administering an effective amount of a compound according to claim 6.

23. A compound of the formula (IA):




(IA)

wherein

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;


n is an integer from 0 to 3;

ZR_1 is the following



wherein

Y_1 is R_3 -(C_1 - C_{12})alkyl, R_4 -aryl, R_5 -heteroaryl, R_6 -(C_3 - C_{12})cyclo-alkyl, R_7 -(C_3 - C_7)heterocycloalkyl, $-CO_2(C_1$ - C_6)alkyl, CN or $-C(O)NR_8R_9$; Y_2 is hydrogen or Y_1 ; Y_3 is hydrogen or (C_1 - C_6)alkyl; or Y_1 , Y_2 and Y_3 , together with the carbon to which they are attached, form one of the following structures:



wherein r is 0 to 3; w and u are each 0-3, provided that the sum of w and u is 1-3; c and d are independently 1 or 2; s is 1 to 5; and ring E is a fused R_4 -phenyl or R_5 -heteroaryl ring;

R_{10} is 1 to 3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-NR_8R_9$ and $-(C_1-C_6)$ alkyl- NR_8R_9 ;


R_{11} is 1 to 3 substituents independently selected from the group consisting of R_{10} , $-CF_3$, $-OCF_3$, NO_2 and halo, or R_{11} substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;

R_8 and R_9 are independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_3-C_{12}) cycloalkyl, aryl and aryl (C_1-C_6) alkyl;

R_3 is 1 to 3 substituents independently selected from the group consisting of H, R_4 -aryl, R_6 -(C_3-C_{12})cycloalkyl, R_5 -heteroaryl, R_7 -(C_3-C_7)heterocycloalkyl, $-NR_8R_9$, $-OR_{12}$ and $-S(O)_{0.2}R_{12}$;

R_6 is 1 to 3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, R_4 -aryl, $-NR_8R_9$, $-OR_{12}$ and $-SR_{12}$;

R_4 is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C_1-C_6) alkyl, R_{13} -aryl, (C_3-C_{12}) cycloalkyl, $-CN$, $-CF_3$, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-OCF_3$, $-NR_8R_9$, $-(C_1-C_6)$ alkyl- NR_8R_9 , $-NHSO_2R_8$, $-SO_2N(R_{14})_2$, $-SO_2R_8$, $-SOR_8$, $-SR_8$, $-NO_2$, $-CONR_8R_9$, $-NR_9COR_8$, $-COR_8$, $-COCF_3$, $-OCOR_8$, $-OCO_2R_8$, $-COOR_8$, $-(C_1-C_6)$ alkyl- $NHCOOC(CH_3)_3$, $-(C_1-C_6)$ alkyl- $NHCOCF_3$, $-(C_1-C_6)$ alkyl- $NHSO_2$ -(C_1-C_6)alkyl, $-(C_1-C_6)$ alkyl- $NHCONH$ -(C_1-C_6)-alkyl and



wherein f is 0 to 6; or R_4 substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;

R_5 is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C_1-C_6) alkyl, R_{13} -aryl, (C_3-C_{12}) cycloalkyl, $-CN$, $-CF_3$, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-OCF_3$, $-NR_8R_9$, $-(C_1-C_6)$ alkyl- NR_8R_9 , $-NHSO_2R_8$, $-SO_2N(R_{14})_2$, $-NO_2$, $-CONR_8R_9$, $-NR_9COR_8$, $-COR_8$, $-OCOR_8$, $-OCO_2R_8$ and $-COOR_8$;

R_7 is H, (C_1-C_6) alkyl, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-NR_8R_9$ or $-(C_1-C_6)$ alkyl- NR_8R_9 ;

R_{12} is H, (C_1-C_6) alkyl, R_4 -aryl, $-(C_1-C_6)$ alkyl- OR_8 , $-(C_1-C_6)$ alkyl- NR_8R_9 , $-(C_1-C_6)$ alkyl- SR_8 , or aryl (C_1-C_6) alkyl;

R_{13} is 1-3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, (C_1-C_6) alkoxy and halo;

R_{14} is independently selected from the group consisting of H, (C_1-C_6) alkyl and R_{13} - $C_6H_4-CH_2-$;

or a pharmaceutically acceptable salt thereof.

24. A pharmaceutical composition comprising a compound of claim 23 and at least one pharmaceutically acceptable excipient.

25. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 23.


26. A method of modulating a pharmacological response from the ORL1 receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 23.

27. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 1.

28. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 6.

29. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 23.

30. A compound of the formula (II):



(II)

wherein

the dotted line represents an optional double bond;

R is hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkylC₁₋₄alkyl-, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy-, C₁₋₁₀ alkenyl, C₁₋₁₀ alkylidene, oxo, C₁₋₁₀ alkyl substituted with 1-3 halogen, C₃₋₁₂ cycloalkyl substituted with 1-3 halogen, C₃₋₁₂ cycloalkylC₁₋₄alkyl- substituted with 1-3 halogen, C₁₋₁₀ alkoxy substituted with 1-3 halogen, C₃₋₁₂ cycloalkoxy- substituted with 1-3 halogen, -COOV₁, -C₁₋₄COOV₁, -CH₂OH, -SO₂N(V₁)₂, hydroxyC₁₋₁₀alkyl-, hydroxyC₃₋₁₀cycloalkyl-, cyanoC₁₋₁₀alkyl-, cyanoC₃₋₁₀cycloalkyl-, -CON(V₁)₂, NH₂SO₂C₁₋₄alkyl-, NH₂SOC₁₋₄alkyl-, sulfonylaminoC₁₋₁₀alkyl-, diaminoalkyl-, -sulfonylC₁₋₄alkyl, a 6-membered heterocyclic ring, a 6-membered heteroaromatic ring, a 6-membered heterocyclicC₁₋₄alkyl-, a 6-membered heteroaromaticC₁₋₄alkyl-, a 6-membered aromatic ring, a 6-membered aromaticC₁₋₄alkyl-, a 5-membered heterocyclic ring optionally substituted with an oxo or thio, a 5-membered heteroaromatic ring, a 5-membered heterocyclicC₁₋₄alkyl- optionally substituted with an oxo or

thio, a 5-membered heteroaromatic C_{1-4} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}(=NH)W_1$, $-C_{1-5}NHC(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, wherein W_1 is hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{1-10} alkoxy, C_{3-12} cycloalkoxy, $-CH_2OH$, amino, C_{1-4} alkylamino-, diC_{1-4} alkylamino-, or a 5-membered heteroaromatic ring optionally substituted with 1-3 lower alkyl;

wherein each V_1 is independently selected from H, C_{1-6} alkyl, C_{3-6} cycloalkyl, benzyl and phenyl;

n is an integer from 0 to 3;


D is a 5-8 membered cycloalkyl, 5-8 membered heterocyclic or a 6 membered aromatic or heteroaromatic group;

n is an integer from 0 to 3;

A , B and Q are independently hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{1-10} alkoxy, C_{3-12} cycloalkoxy, C_{1-10} alkenyl, C_{1-10} alkylidene, oxo, $-CH_2OH$, $-NHSO_2$, hydroxy C_{1-10} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, diC_{1-4} alkylaminocarbonyl-, acylamino-, acylaminoalkyl-, amide, sulfonylamino C_{1-10} alkyl-, or A - B can together form a C_{2-6} bridge, or B - Q can together form a C_{3-7} bridge, or A - Q can together form a C_{1-5} bridge;

Z is selected from the group consisting of a bond, straight or branched C_{1-6} alkylene, $-NH-$, $-CH_2O-$, $-CH_2NH-$, $-CH_2N(CH_3)-$, $-NHCH_2-$, $-CH_2CONH-$, $-NHCH_2CO-$, $-CH_2CO-$, $-COCH_2-$, $-CH_2COCH_2-$, $-CH(CH_3)-$, $-CH=$, $-O-$ and $-HC=CH-$, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with one or more lower alkyl, hydroxy, halo or alkoxy group;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, $-COOV_1$, $-C_{1-4}COOV_1$, cyano, cyano C_{1-10} alkyl-, cyano C_{3-10} cycloalkyl-, NH_2SO_2- , $NH_2SO_2C_{1-4}$ alkyl-, NH_2SOC_{1-4} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, diC_{1-4} alkylaminocarbonyl-, benzyl, C_{3-12} cycloalkenyl-, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a hetero-monocyclic ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ; and wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, or benzyl of R_1 is optionally substituted with 1-3 substituents selected from the group consisting of halogen, hydroxy, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, cyano-, $-COOV_1$, $-C_{1-4}COOV_1$, cyano C_{1-10} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, a 5-membered heteroaromatic C_{0-4} alkyl-, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl-, C_{1-10} alkoxy-, and cyano; and wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, or spiro ring system of the formula (V) is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy or benzyloxy is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl-and halogen, said alkyl or cycloalkyl optionally substituted with an oxo, amino, alkylamino or dialkylamino group;

or a pharmaceutically acceptable salt thereof or solvate thereof.

31. A compound of claim 30, wherein D is phenyl or a 6 membered heteroaromatic group containing 1-3 nitrogen atoms.

32. A compound of claim 30, wherein R is selected from the group consisting of $-CH_2C=ONH_2$, $-C(NH)NH_2$, pyridylmethyl, cyclopentyl, cyclohexyl, furanylmethyl, $-C=OCH_3$, $-CH_2CH_2NHC=OCH_3$, $-SO_2CH_3$, $CH_2CH_2NHSO_2CH_3$, furanylcabonyl-, methylpyrrolylcabonyl-, diazolecarbonyl-, azolemethyl-, trifluoroethyl-, hydroxyethyl-, cyanomethyl-, oxo-oxazolemethyl-, and diazolemethyl-.


33. A compound of claim 30, wherein ZR_1 is selected from the group consisting of cyclohexylethyl-, cyclohexylmethyl-, cyclopentylmethyl-, dimethylcyclohexylmethyl-, phenylethyl-, pyrrolyltrifluoroethyl-, thienyltrifluoroethyl-, pyridylethyl-, cyclopentyl-, cyclohexyl-, methoxycyclohexyl-, tetrahydropyranyl-, propylpiperidinyl-, indolylmethyl-, pyrazoylpentyl-, thiazolyethyl-, phenyltrifluoroethyl-, hydroxyhexyl-, methoxyhexyl-,

isopropoxybutyl-, hexyl-, and oxocanylpropyl-.

34. A compound of claim 30, wherein at least one of ZR_1 or R is selected from the group consisting of CH_2COOV_1 , tetrazolylmethyl-, cyanomethyl-, NH_2SO_2 methyl-, NH_2SO methyl-, aminocarbonylmethyl-, C_{1-4} alkylaminocarbonylmethyl-, and diC_{1-4} alkylaminocarbonylmethyl-.

35. A compound of claim 30, wherein ZR_1 is 3,3 diphenylpropyl optionally substituted at the 3 carbon of the propyl with $-COOV_1$, tetrazolyl C_{0-4} alkyl-, cyano-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, or diC_{1-4} alkylaminocarbonyl-.

36. A compound of the formula (IIA):



(IIA)

wherein


the dotted line represents an optional double bond;

Z is selected from the group consisting of a bond, $-CH_2-$, $-NH-$, $-CH_2O-$, $-CH_2CH_2-$, $-CH_2NH-$, $-CH_2N(CH_3)-$, $-NHCH_2-$, $-CH_2CONH-$, $-NHCH_2CO-$, $-CH_2CO-$, $-COCH_2-$, $-CH_2COCH_2-$, $-CH(CH_3)-$, $-CH=$, and $-HC=CH-$, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with a lower alkyl, halogen, hydroxy or alkoxy group;

R and Q are the same or different and are each selected from the group consisting of

hydrogen, halogen, C_{1-10} alkyl, C_{1-10} alkenyl, C_{1-10} alkylidene, C_{3-12} cycloalkyl, C_{1-10} alkoxy, and oxo;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, benzyl, C_{3-12} cycloalkenyl, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a heteromonocyclic ring, a bicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ;

wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, or benzyl is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, cyano, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, heteromonocyclic ring, heterobicyclic ring system, and spiro ring system of the formula (V) are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy and benzyloxy are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

or a pharmaceutically acceptable salt thereof.

37. A compound of claim 36, wherein Q is hydrogen or methyl.

38. A compound of claim 36, wherein R is hydrogen, methyl, ethyl, or ethylidene.
39. A compound of claim 36, wherein R₁ is alkyl selected from the group consisting of methyl, ethyl, propyl, butyl, pentyl and hexyl.
40. A compound of claim 36, wherein R₁ is cycloalkyl selected from the group consisting of cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, and norbornyl.
41. A compound of claim 36, wherein R₁ is tetrahydronaphthyl, decahydronaphthyl or dibenzocycloheptyl.
42. A compound of claim 36, wherein R₁ is phenyl or benzyl.
43. A compound of claim 36, wherein R₁ is a bicyclic aromatic ring.
44. A compound of claim 43, wherein said bicyclic aromatic ring is indenyl, quinoline or naphthyl.
45. A compound of claim 36, wherein Z is a bond, methyl, or ethyl.
46. A compound of claim 36, wherein n is 0.
47. A compound of claim 36, wherein X₁ and X₂ are both O.
48. A compound of claim 36, wherein the dotted line is a double bond.
49. A compound selected from the group consisting of:
3-ethylidene-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
3-ethylidene-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
3-ethylidene-1-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
3-ethylidene-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
3-ethylidene-1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
3-ethylidene-1-[1-(*p*-benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
3-ethylidene-1-[1-(benzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethylidene-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethylidene-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethylidene-1-[1-(3,3-diphenylpropyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethylidene-1-[1-(*p*-cyanobenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethyl-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethyl-1-[1-[4-(1-methylethyl)-cyclohexyl]-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethyl-1-[1-(4-propylcyclohexyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethyl-1-[1-(1,2,3,4-tetrahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethyl-1-[1-(decahydro-2-naphthyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethyl-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethyl-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(naphth-1-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(naphth-2-yl-methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(*p*-phenylbenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(3,3-Bis(phenyl)propyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(*p*-cyanobenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(*p*-benzyloxybenzyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(1,2,3,4-tetrahydronaphth-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(benzyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(4-propyl-cyclohexyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(5-methylhex-2-yl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(decahydro-2-naphthyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(4-(1-methylethyl)-cyclohexyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(cyclooctylmethyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 1-[1-(3,3-Bis(phenyl)propyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;
 3-ethyl-1-[1-(3,3-Bis(phenyl)propyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(4-propylcyclohexyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(5-methylhex-2-yl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-[4-(1-methylethyl)cyclohexyl]-3-methyl-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

3-ethyl-1-[1-(decahydro-2-naphthyl)-3-(methyl)-4-piperidinyl]-1,3-dihydro-2H-indole-2-one;

and pharmaceutically acceptable salts thereof.

50. A pharmaceutical composition comprising a compound of claim 30 and at least one pharmaceutically acceptable excipient.

51. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 30.


52. A method of modulating a pharmacological response from the ORL1 receptor comprising administering to a patient in need thereof an effective amount of a compound according to claim 30.

53. A pharmaceutical composition comprising a compound of claim 36 and at least one pharmaceutically acceptable excipient.

54. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 36.

55. A method of modulating a pharmacological response from the ORL1 receptor comprising administering an effective amount of a compound according to claim 36.

56. A compound of the formula (IIA):



(IIA)


wherein

the dotted line represents an optional double bond;

R and Q are the same or different and are each selected from the group consisting of hydrogen, halogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkenyl, C₁₋₁₀ alkylidene, C₃₋₁₂ cycloalkyl, C₁₋₁₀ alkoxy, and oxo;





R₂ is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

ZR₁ is the following




wherein

Y_1 is R_3 -(C_1 - C_{12})alkyl, R_4 -aryl, R_5 -heteroaryl, R_6 -(C_3 - C_{12})cyclo-alkyl, R_7 -(C_3 - C_7)heterocycloalkyl, $-CO_2(C_1$ - C_6)alkyl, CN or $-C(O)NR_8R_9$; Y_2 is hydrogen or Y_1 ; Y_3 is hydrogen or (C_1 - C_6)alkyl; or Y_1 , Y_2 and Y_3 , together with the carbon to which they are attached, form one of the following structures:



or



wherein r is 0 to 3; w and u are each 0-3, provided that the sum of w and u is 1-3; c and d are independently 1 or 2; s is 1 to 5; and ring E is a fused R₄-phenyl or R₅-heteroaryl ring;

R₁₀ is 1 to 3 substituents independently selected from the group consisting of H, (C₁-C₆)alkyl, -OR₈, -(C₁-C₆)alkyl-OR₈, -NR₈R₉ and -(C₁-C₆)alkyl-NR₈R₉;


R₁₁ is 1 to 3 substituents independently selected from the group consisting of R₁₀, -CF₃, -OCF₃, NO₂ and halo, or R₁₁ substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;

R₈ and R₉ are independently selected from the group consisting of hydrogen, (C₁-C₆)alkyl, (C₃-C₁₂)cycloalkyl, aryl and aryl(C₁-C₆)alkyl;

R₃ is 1 to 3 substituents independently selected from the group consisting of H, R₄-aryl, R₆-(C₃-C₁₂)cycloalkyl, R₅-heteroaryl, R₇-(C₃-C₇)heterocycloalkyl, -NR₈R₉, -OR₁₂ and -S(O)₀₋₂R₁₂;

R₆ is 1 to 3 substituents independently selected from the group consisting of H, (C₁-C₆)alkyl, R₄-aryl, -NR₈R₉, -OR₁₂ and -SR₁₂;

R₄ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C₁-C₆)alkyl, R₁₃-aryl, (C₃-C₁₂)cycloalkyl, -CN, -CF₃, -OR₈, -(C₁-C₆)alkyl-OR₈, -OCF₃, -NR₈R₉, -(C₁-C₆)alkyl-NR₈R₉, -NHSO₂R₈, -SO₂N(R₁₄)₂, -SO₂R₈, -SOR₈, -SR₈, -NO₂, -CONR₈R₉, -NR₉COR₈, -COR₈, -COCF₃, -OCOR₈, -OCO₂R₈, -COOR₈, -(C₁-C₆)alkyl-NHCOOC(CH₃)₃, -(C₁-C₆)alkyl-NHCOCF₃, -(C₁-C₆)alkyl-NHSO₂-(C₁-C₆)alkyl, -(C₁-C₆)alkyl-NHCONH-(C₁-C₆)alkyl and



wherein f is 0 to 6; or R₄ substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;

R₅ is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C₁-C₆)alkyl, R₁₃-aryl, (C₃-C₁₂)cycloalkyl, -CN, -CF₃, -OR₈, -(C₁-C₆)alkyl-OR₈, -OCF₃, -NR₈R₉, -(C₁-C₆)alkyl-NR₈R₉, -NHSO₂R₈, -SO₂N(R₁₄)₂, -NO₂, -CONR₈R₉, -NR₉COR₈, -COR₈, -OCOR₈, -OCO₂R₈ and -COOR₈;

R₇ is H, (C₁-C₆)alkyl, -OR₈, -(C₁-C₆)alkyl-OR₈, -NR₈R₉ or -(C₁-C₆)alkyl-NR₈R₉;

R₁₂ is H, (C₁-C₆)alkyl, R₄-aryl, -(C₁-C₆)alkyl-OR₈, -(C₁-C₆)alkyl-NR₈R₉, -(C₁-C₆)alkyl-SR₈, or aryl (C₁-C₆)alkyl;

R_{13} is 1-3 substituents independently selected from the group consisting of H, (C₁-C₆)alkyl, (C₁-C₆)alkoxy and halo;

R_{14} is independently selected from the group consisting of H, (C₁-C₆)alkyl and R_{13} -C₆H₄-CH₂-;

or a pharmaceutically acceptable salt thereof.

57. A pharmaceutical composition comprising a compound of claim 56 and at least one pharmaceutically acceptable excipient.

58. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 56.


59. A method of modulating a pharmacological response from the ORL1 receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 56.

60. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 30.

61. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 36.

62. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 56.

63. A compound of the (III):



(III)

wherein R is hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkylC₁₋₄alkyl-, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy-, C₁₋₁₀ alkyl substituted with 1-3 halogen, C₃₋₁₂ cycloalkyl substituted with 1-3 halogen, C₃₋₁₂ cycloalkylC₁₋₄alkyl- substituted with 1-3 halogen, C₁₋₁₀ alkoxy substituted with 1-3 halogen, C₃₋₁₂ cycloalkoxy- substituted with 1-3 halogen, -COOV₁, -C₁₋₄COOV₁, -CH₂OH, -SO₂N(V₁)₂, hydroxyC₁₋₁₀alkyl-, hydroxyC₃₋₁₀cycloalkyl-, cyanoC₁₋₁₀alkyl-, cyanoC₃₋₁₀cycloalkyl-, -CON(V₁)₂, NH₂SO₂C₁₋₄alkyl-, NH₂SOC₁₋₄alkyl-, sulfonylaminoC₁₋₁₀alkyl-, diaminoalkyl-, -sulfonylC₁₋₄alkyl, a 6-membered heterocyclic ring, a 6-membered heteroaromatic ring, a 6-membered heterocyclicC₁₋₄alkyl-, a 6-membered heteroaromaticC₁₋₄alkyl-, a 6-membered aromatic ring, a 6-membered aromaticC₁₋₄alkyl-, a 5-membered heterocyclic ring optionally substituted with an oxo or thio, a 5-membered heteroaromatic ring, a 5-membered heterocyclicC₁₋₄alkyl- optionally substituted with an oxo or thio, a 5-membered heteroaromaticC₁₋₄alkyl-, -C₁₋₅(=O)W₁, -C₁₋₅(=NH)W₁, -C₁₋₅NHC(=O)W₁, -C₁₋₅NHS(=O)₂W₁, -C₁₋₅NHS(=O)W₁, wherein W₁ is hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy, -CH₂OH, amino, C₁₋₄alkylamino-, diC₁₋₄alkylamino-, or a 5-membered heteroaromatic ring optionally substituted with 1-3 lower alkyl;


wherein each V₁ is independently selected from H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, benzyl and phenyl;

n is an integer from 0 to 3;

D is a 5-8 membered cycloalkyl, 5-8 membered heterocyclic or a 6 membered aromatic

or heteroaromatic group;


Z is selected from the group consisting of a bond, straight or branched C_{1-6} alkylene, -NH-, $-CH_2O-$, $-CH_2NH-$, $-CH_2N(CH_3)-$, $-NHCH_2-$, $-CH_2CONH-$, $-NHCH_2CO-$, $-CH_2CO-$, $-COCH_2-$, $-CH_2COCH_2-$, $-CH(CH_3)-$, $-CH=$, $-O-$ and $-HC=CH-$, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with one or more lower alkyl, hydroxy, halo or alkoxy group; or Z is a cycloalkylamino system of the formula (VI):



(VI)

wherein A, B and Q are independently hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{1-10} alkoxy, C_{3-12} cycloalkoxy, $-CH_2OH$, $-NHSO_2$, hydroxy C_{1-10} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, di C_{1-4} alkylaminocarbonyl-, acylamino-, acylaminoalkyl-, amide, sulfonylamino C_{1-10} alkyl-, or A-B can together form a C_{2-6} bridge, or B-Q can together form a C_{3-7} bridge, or A-Q can together form a C_{1-5} bridge;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, $-COOV_1$, $-C_{1-4}COOV_1$, cyano, cyano C_{1-10} alkyl-, cyano C_{3-10} cycloalkyl-, NH_2SO_2 -, $NH_2SO_2C_{1-4}$ alkyl-, NH_2SOC_{1-4} alkyl-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, di C_{1-4} alkylaminocarbonyl-, benzyl, C_{3-12} cycloalkenyl-, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a hetero-monocyclic ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ; and wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, or benzyl of R_1 is optionally substituted with 1-3 substituents selected from the group consisting of halogen, hydroxy, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, cyano, $-COOV_1$, $-C_{1-4}COOV_1$, cyano C_{1-10} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, a 5-membered heteroaromatic C_{0-4} alkyl-, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl-, C_{1-10} alkoxy-, and cyano; and wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, or spiro ring system of the formula (V) is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy or benzyloxy is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl-and halogen, said alkyl or cycloalkyl optionally substituted with an oxo, amino, alkylamino or dialkylamino group;

or a pharmaceutically acceptable salt thereof or solvate thereof.


64. A compound of claim 63, wherein D is phenyl or a 6 membered heteroaromatic group containing 1-3 nitrogen atoms.

65. A compound of claim 63, wherein R is selected from the group consisting of $-CH_2C=ONH_2$, $-C(NH)NH_2$, pyridylmethyl, cyclopentyl, cyclohexyl, furanylmethyl, $-C=OCH_3$, $-CH_2CH_2NHC=OCH_3$, $-SO_2CH_3$, $CH_2CH_2NHSO_2CH_3$, furanylcarbonyl-, methylpyrrolylcarbonyl-, diazolecarbonyl-, azolemethyl-, trifluoroethyl-, hydroxyethyl-, cyanomethyl-, oxo-oxazolemethyl-, and diazolemethyl-.

66. A compound of claim 63, wherein ZR_1 is selected from the group consisting of cyclohexylethyl-, cyclohexylmethyl-, cyclopentylmethyl-, dimethylcyclohexylmethyl-, phenylethyl-, pyrrolyltrifluoroethyl-, thienyltrifluoroethyl-, pyridylethyl-, cyclopentyl-, cyclohexyl-, methoxycyclohexyl-, tetrahydropyranyl-, propylpiperidinyl-, indolylmethyl-, pyrazolypentyl-, thiazolylethyl-, phenyltrifluoroethyl-, hydroxyhexyl-, methoxyhexyl-, isopropoxybutyl-, hexyl-, and oxocanylpropyl-.

67. A compound of claim 63, wherein at least one of ZR_1 or R is selected from the group consisting of CH_2COOV_1 , tetrazolylmethyl-, cyanomethyl-, NH_2SO_2 methyl-, NH_2SO methyl-, aminocarbonylmethyl-, C_{1-4} alkylaminocarbonylmethyl-, and diC_{1-4} alkylaminocarbonylmethyl-.
68. A compound of claim 63, wherein ZR_1 is 3,3 diphenylpropyl optionally substituted at the 3 carbon of the propyl with $-COOV_1$, tetrazolyl C_{0-4} alkyl-, cyano-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, or diC_{1-4} alkylaminocarbonyl-.

69. A compound of formula (IIIA):




(IIIA)

wherein

n is an integer from 0 to 3;

Z is selected from the group consisting of a bond, $-CH_2-$, $-NH-$, $-CH_2O-$, $-CH_2CH_2-$, $-CH_2NH-$, $-CH_2N(CH_3)-$, $-NHCH_2-$, $-CH_2CONH-$, $-NHCH_2CO-$, $-CH_2CO-$, $-COCH_2-$, $-CH_2COCH_2-$, $-CH(CH_3)-$, $-CH=$, $-HC=CH-$, and a cycloalkylamino system of the formula (VI):




(VI)

wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with a lower alkyl, halogen, hydroxy, phenyl, benzyl, or alkoxy group;

R is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{1-10} alkoxy, and C_{3-12} cycloalkyl;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, benzyl, C_{3-12} cycloalkenyl, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a heteromonocyclic ring, a heterobicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ;

wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, or benzyl is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, cyano, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, heteromonocyclic ring, heterobicyclic ring system, and spiro ring system of the formula (V) are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl, phenyl, benzyl, phenyloxy and

benzyloxy, wherein said phenyl, benzyl, phenyloxy and benzyloxy are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, and cyano;

R₂ is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;
or a pharmaceutically acceptable salt thereof.

70. A compound of claim 69, wherein R₁ is alkyl selected from the group consisting of methyl, ethyl, propyl, butyl, pentyl and hexyl.
71. A compound of claim 69, wherein R₁ is cycloalkyl selected from the group consisting of cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, and norbornyl.
72. A compound of claim 69, wherein R₁ is tetrahydronaphthyl, decahydronaphthyl or dibenzocycloheptyl.
73. A compound of claim 69, wherein R₁ is phenyl or benzyl.
74. A compound of claim 69, wherein R₁ is a bicyclic aromatic ring.
75. A compound of claim 74, wherein said bicyclic aromatic ring is indenyl, quinoline or naphthyl.
76. A compound of claim 69, wherein Z is a bond, methyl, or ethyl.
77. A compound of claim 69, wherein n is 0.
78. A compound of claim 69, wherein X₁ and X₂ are both O.
79. A compound of claim 69, wherein Z is a cycloalkylamino system of the formula (VI).
80. A compound of claim selected from the group consisting of
3-ethyl-1-(*p*-phenylbenzyl)-1,3-dihydro-2H-benzimidazol-2-one;
3-ethyl-1-(5-methylhex-2-yl)-1,3-dihydro-2H-benzimidazol-2-one;
3-ethyl-1-(4-propylcyclohexyl)-1,3-dihydro-2H-benzimidazol-2-one;

3-ethyl-1-(decahydro-2-naphthyl)-1,3-dihydro-2H-benzimidazol-2-one;
3-ethyl-1-(naphth-2-yl-methyl)-1,3-dihydro-2H-benzimidazol-2-one;
1-(*p*-benzyloxybenzyl)-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-benzyl-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(benzylamino)-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
3-ethyl-1-(naphthylmethyl)-1,3-dihydro-2H-benzimidazol-2-one;
3-ethyl-1-[5-(3-fluorophenyl)-5-(4-fluorophenyl)-hexyl]-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(naphth-2-yl-methyl)ethylamino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(norbornan-2-ylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[[4-(1-methylethyl)-cyclohexyl]amino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(ethylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(benzylamino)-cyclohexyl]-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(indan-2-yl)benzylamino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(cyclooctylmethyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(naphth-2-yl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(*p*-benzyloxybenzyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(cyclooctylmethyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-3-ethyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(benzylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-(dibenzylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(*p*-phenylbenzyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(1,2,3,4-tetrahydronaphthyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;
1-[4-[(4-propyl-cyclohexyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-[(5-methylhex-2-yl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-[(decahydro-2-naphthyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-(cyclooctylamino)-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-[(indan-2-yl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-[(4-phenyl-cyclohexyl)amino]-cyclohexyl]-5-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one;

1-[4-[(5-methylhex-2-yl)amino]-cyclohexyl]-7-carbamoyl-1,3-dihydro-2H-benzimidazol-2-one; and

pharmaceutically acceptable salts thereof.

81. A pharmaceutical composition comprising a compound of claim 63 and at least one pharmaceutically acceptable excipient.

82. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 63.


83. A method of modulating a pharmacological response from the ORL1 receptor comprising administering to a patient in need thereof an effective amount of a compound according to claim 63.

84. A pharmaceutical composition comprising a compound of claim 69 and at least one pharmaceutically acceptable excipient.

85. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 69.

86. A method of modulating a pharmacological response from the ORL1 receptor comprising administering an effective amount of a compound according to claim 69.

87. A compound of claim (IIIA):



(IIIA)


wherein

n is an integer from 0 to 3;

R is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, and C₃₋₁₂ cycloalkyl;


R₂ is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

ZR₁ is the following



wherein

Y₁ is R₃-(C₁-C₁₂)alkyl, R₄-aryl, R₅-heteroaryl, R₆-(C₃-C₁₂)cyclo-alkyl, R₇-(C₃-C₇)heterocycloalkyl, -CO₂(C₁-C₆)alkyl, CN or -C(O)NR₈R₉; Y₂ is hydrogen or Y₁; Y₃ is hydrogen or (C₁-C₆)alkyl; or Y₁, Y₂ and Y₃, together with the carbon to which they are attached, form one of the following structures:



or

wherein r is 0 to 3; w and u are each 0-3, provided that the sum of w and u is 1-3; c and d are independently 1 or 2; s is 1 to 5; and ring E is a fused R_4 -phenyl or R_5 -heteroaryl ring;

R_{10} is 1 to 3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, $-OR_8$, $-(C_1-C_6)alkyl-OR_8$, $-NR_8R_9$ and $-(C_1-C_6)alkyl-NR_8R_9$;

R_{11} is 1 to 3 substituents independently selected from the group consisting of R_{10} , $-CF_3$, $-OCF_3$, NO_2 and halo, or R_{11} substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;


R_8 and R_9 are independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_3-C_{12}) cycloalkyl, aryl and aryl $(C_1-C_6)alkyl$;

R_3 is 1 to 3 substituents independently selected from the group consisting of H, R_4 -aryl, R_6 -(C_3-C_{12})cycloalkyl, R_5 -heteroaryl, R_7 -(C_3-C_7)heterocycloalkyl, $-NR_8R_9$, $-OR_{12}$ and -

$S(O)_{0.2}R_{12}$;

R_6 is 1 to 3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, R_4 -aryl, $-NR_8R_9$, $-OR_{12}$ and $-SR_{12}$;

R_4 is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C_1-C_6) alkyl, R_{13} -aryl, (C_3-C_{12}) cycloalkyl, $-CN$, $-CF_3$, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-OCF_3$, $-NR_8R_9$, $-(C_1-C_6)$ alkyl- NR_8R_9 , $-NHSO_2R_8$, $-SO_2N(R_{14})_2$, $-SO_2R_8$, $-SOR_8$, $-SR_8$, $-NO_2$, $-CONR_8R_9$, $-NR_9COR_8$, $-COR_8$, $-COCF_3$, $-OCOR_8$, $-OCO_2R_8$, $-COOR_8$, $-(C_1-C_6)$ alkyl- $NHCOOC(CH_3)_3$, $-(C_1-C_6)$ alkyl- $NHCOCF_3$, $-(C_1-C_6)$ alkyl- $NHSO_2-(C_1-C_6)$ alkyl, $-(C_1-C_6)$ alkyl- $NHCONH-(C_1-C_6)$ alkyl and



wherein f is 0 to 6; or R_4 substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;

R_5 is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C_1-C_6) alkyl, R_{13} -aryl, (C_3-C_{12}) cycloalkyl, $-CN$, $-CF_3$, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-OCF_3$, $-NR_8R_9$, $-(C_1-C_6)$ alkyl- NR_8R_9 , $-NHSO_2R_8$, $-SO_2N(R_{14})_2$, $-NO_2$, $-CONR_8R_9$, $-NR_9COR_8$, $-COR_8$, $-OCOR_8$, $-OCO_2R_8$ and $-COOR_8$;

R_7 is H, (C_1-C_6) alkyl, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-NR_8R_9$ or $-(C_1-C_6)$ alkyl- NR_8R_9 ;

R_{12} is H, (C_1-C_6) alkyl, R_4 -aryl, $-(C_1-C_6)$ alkyl- OR_8 , $-(C_1-C_6)$ alkyl- NR_8R_9 , $-(C_1-C_6)$ alkyl- SR_8 , or aryl (C_1-C_6) alkyl;

R_{13} is 1-3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, (C_1-C_6) alkoxy and halo;

R_{14} is independently selected from the group consisting of H, (C_1-C_6) alkyl and $R_{13}-C_6H_4-CH_2-$;

or a pharmaceutically acceptable salt thereof.

88. A pharmaceutical composition comprising a compound of claim 87 and at least one pharmaceutically acceptable excipient.

89. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 87.


90. A method of modulating a pharmacological response from the ORL1 receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 87.

91. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 63.

92. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 69.

93. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 87.

94. A compound of formula (IV):



(IV)

wherein R is hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkylC₁₋₄alkyl-, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy-, C₁₋₁₀ alkyl substituted with 1-3 halogen, C₃₋₁₂ cycloalkyl substituted with 1-3 halogen, C₃₋₁₂ cycloalkylC₁₋₄alkyl- substituted with 1-3 halogen, C₁₋₁₀ alkoxy substituted with 1-3 halogen, C₃₋₁₂ cycloalkoxy- substituted with 1-3 halogen, -COOV₁, -C₁₋₄COOV₁, -CH₂OH, -SO₂N(V₁)₂, hydroxyC₁₋₁₀alkyl-, hydroxyC₃₋₁₀cycloalkyl-, cyanoC₁₋₁₀alkyl-, cyanoC₃₋₁₀cycloalkyl-, -CON(V₁)₂, NH₂SO₂C₁₋₄alkyl-, NH₂SOC₁₋₄alkyl-, sulfonylaminoC₁₋₁₀alkyl-, diaminoalkyl-, -sulfonylC₁₋₄alkyl, a 6-membered heterocyclic ring, a 6-membered heteroaromatic ring, a 6-membered heterocyclicC₁₋₄alkyl-, a 6-membered heteroaromaticC₁₋₄alkyl-, a 6-membered aromatic ring, a 6-membered aromaticC₁₋₄alkyl-, a 5-membered heterocyclic ring optionally substituted with an oxo or thio, a 5-membered heteroaromatic ring, a 5-membered heterocyclicC₁₋₄alkyl- optionally substituted with an oxo or thio, a 5-membered heteroaromaticC₁₋₄alkyl-, -C₁₋₅(=O)W₁, -C₁₋₅(=NH)W₁, -C₁₋₅NHC(=O)W₁, -C₁₋₅NHS(=O)₂W₁, -C₁₋₅NHS(=O)W₁, wherein W₁ is hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy, -CH₂OH, amino, C₁₋₄alkylamino-, diC₁₋₄alkylamino-, or a 5-membered heteroaromatic ring optionally substituted with 1-3 lower alkyl;

wherein each V₁ is independently selected from H, C₁₋₆ alkyl, C₃₋₆ cycloalkyl, benzyl and phenyl;

D is a 5-8 membered cycloalkyl, 5-8 membered heterocyclic or a 6 membered aromatic or heteroaromatic group;


n is an integer from 0 to 3;

A, B and Q are independently hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl, C₁₋₁₀ alkoxy, C₃₋₁₂ cycloalkoxy, -CH₂OH, -NHSO₂, hydroxyC₁₋₁₀alkyl-, aminocarbonyl-, C₁₋₄alkylaminocarbonyl-, diC₁₋₄alkylaminocarbonyl-, acylamino-, acylaminoalkyl-, amide, sulfonylaminoC₁₋₁₀alkyl-, or A-B can together form a C₂₋₆ bridge, or B-Q can together form a C₃₋₇ bridge, or A-Q can together form a C₁₋₅ bridge;

Z is selected from the group consisting of a bond, straight or branched C₁₋₆ alkylene, -NH-, -CH₂O-, -CH₂NH-, -CH₂N(CH₃)-, -NHCH₂-, -CH₂CONH-, -NHCH₂CO-, -CH₂CO-, -COCH₂-, -CH₂COCH₂-, -CH(CH₃)-, -CH=, -O- and -HC=CH-, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with one or more lower alkyl, hydroxy, halo or alkoxy group;

R₁ is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂cycloalkyl, C₂₋₁₀alkenyl, amino, C₁₋₁₀alkylamino-, C₃₋₁₂cycloalkylamino-, -COOV₁, -C₁₋₄COOV₁, cyano, cyanoC₁₋₁₀alkyl-, cyanoC₃₋₁₀cycloalkyl-, NH₂SO₂-, NH₂SO₂C₁₋₄alkyl-, NH₂SOC₁₋₄alkyl-, aminocarbonyl-, C₁₋₄alkylaminocarbonyl-, diC₁₋₄alkylaminocarbonyl-, benzyl, C₃₋₁₂ cycloalkenyl-, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a hetero-monocyclic

ring, a hetero-bicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X_1 and X_2 are independently selected from the group consisting of NH, O, S and CH_2 ; and wherein said alkyl, cycloalkyl, alkenyl, C_{1-10} alkylamino-, C_{3-12} cycloalkylamino-, or benzyl of R_1 is optionally substituted with 1-3 substituents selected from the group consisting of halogen, hydroxy, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, cyano-, $COOV_1$, $-C_{1-4}COOV_1$, cyano C_{1-10} alkyl-, $-C_{1-5}(=O)W_1$, $-C_{1-5}NHS(=O)_2W_1$, $-C_{1-5}NHS(=O)W_1$, a 5-membered heteroaromatic C_{0-4} alkyl-, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl-, C_{1-10} alkoxy-, and cyano; and wherein said C_{3-12} cycloalkyl, C_{3-12} cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, hetero-monocyclic ring, hetero-bicyclic ring system, or spiro ring system of the formula (II) is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, nitro, trifluoromethyl-, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy or benzyloxy is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C_{1-10} alkyl, C_{1-10} alkoxy, and cyano;

R_2 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl- and halogen, said alkyl or cycloalkyl optionally substituted with an oxo, amino, alkylamino or dialkylamino group;

or a pharmaceutically acceptable salt thereof or solvate thereof.

95. A compound of claim 94, wherein D is phenyl or a 6 membered heteroaromatic group containing 1-3 nitrogen atoms.

96. A compound of claim 94, wherein R is selected from the group consisting of $-CH_2C=ONH_2$, $-C(NH)NH_2$, pyridylmethyl, cyclopentyl, cyclohexyl, furanylmethyl, $-C=OCH_3$, $-CH_2CH_2NHC=OCH_3$, $-SO_2CH_3$, $CH_2CH_2NHSO_2CH_3$, furanylcarbonyl-, methylpyrrolylcarbonyl-, diazolecarbonyl-, azolemethyl-, trifluoroethyl-, hydroxyethyl-,


cyanomethyl-, oxo-oxazolomethyl-, and diazomethyl-.

97. A compound of claim 94, wherein ZR_1 is selected from the group consisting of cyclohexylethyl-, cyclohexylmethyl-, cyclopentylmethyl-, dimethylcyclohexylmethyl-, phenylethyl-, pyrrolyltrifluoroethyl-, thienyltrifluoroethyl-, pyridylethyl-, cyclopentyl-, cyclohexyl-, methoxycyclohexyl-, tetrahydropyranyl-, propylpiperidinyl-, indolylmethyl-, pyrazolypentyl-, thiazolylethyl-, phenyltrifluoroethyl-, hydroxyhexyl-, methoxyhexyl-, isopropoxybutyl-, hexyl-, and oxocanylpropyl-.

98. A compound of claim 94, wherein at least one of ZR_1 or R is selected from the group consisting of CH_2COOV_1 , tetrazolylmethyl-, cyanomethyl-, NH_2SO_2 methyl-, NH_2SO methyl-, aminocarbonylmethyl-, C_{1-4} alkylaminocarbonylmethyl-, and diC_{1-4} alkylaminocarbonylmethyl-.

99. A compound of claim 94, wherein ZR_1 is 3,3 diphenylpropyl optionally substituted at the 3 carbon of the propyl with $-COOV_1$, tetrazolyl C_{0-4} alkyl-, cyano-, aminocarbonyl-, C_{1-4} alkylaminocarbonyl-, or diC_{1-4} alkylaminocarbonyl-.

100.. A compound of the formula (IVA):



(IVA)


wherein

n is an integer from 0 to 3;

Z is selected from the group consisting of a bond, $-CH_2-$, $-NH-$, $-CH_2O-$, $-CH_2CH_2-$, $-CH_2NH-$, $-CH_2N(CH_3)-$, $-NHCH_2-$, $-CH_2CONH-$, $-NHCH_2CO-$, $-CH_2CO-$, $-COCH_2-$, $-CH_2COCH_2-$, $-CH(CH_3)-$, $-CH=$, and $-HC=CH-$, wherein the carbon and/or nitrogen atoms are unsubstituted or substituted with a lower alkyl, halogen, hydroxy or alkoxy group;

R is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{1-10} alkoxy, and C_{3-12} cycloalkyl;

R_1 is selected from the group consisting of hydrogen, C_{1-10} alkyl, C_{3-12} cycloalkyl, C_{2-10} alkenyl, amino, C_{1-10} alkylamino, C_{3-12} cycloalkylamino, benzyl, C_{3-12} cycloalkenyl, a monocyclic, bicyclic or tricyclic aryl or heteroaryl ring, a heteromonocyclic ring, a heterobicyclic ring system, and a spiro ring system of the formula (V):



(V)

wherein X₁ and X₂ are independently selected from the group consisting of NH, O, S and CH₂;

wherein said alkyl, cycloalkyl, alkenyl, C₁₋₁₀alkylamino, C₃₋₁₂cycloalkylamino, or benzyl is optionally substituted with 1-3 substituents selected from the group consisting of halogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, nitro, trifluoromethyl, cyano, phenyl, benzyl, benzyloxy, said phenyl, benzyl, and benzyloxy optionally being substituted with 1-3 substituents selected from the group consisting of halogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, and cyano;

wherein said C₃₋₁₂ cycloalkyl, C₃₋₁₂ cycloalkenyl, monocyclic, bicyclic or tricyclic aryl, heteroaryl ring, heteromonocyclic ring, heterobicyclic ring system, and spiro ring system of the formula (V) are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, nitro, trifluoromethyl, phenyl, benzyl, phenyloxy and benzyloxy, wherein said phenyl, benzyl, phenyloxy and benzyloxy are optionally substituted with 1-3 substituents selected from the group consisting of halogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, and cyano;

R₂ is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

or a pharmaceutically acceptable salt thereof.

101. A compound of claim 100, wherein R₁ is alkyl selected from the group consisting of methyl, ethyl, propyl, butyl, pentyl and hexyl.

102. A compound of claim 100, wherein R₁ is cycloalkyl selected from the group consisting of cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, and norbornyl.

103. A compound of claim 100, wherein R₁ is tetrahydronaphthyl, decahydronaphthyl or dibenzocycloheptyl.

104. A compound of claim 100, wherein R_1 is phenyl or benzyl.
105. A compound of claim 100, wherein R_1 is a bicyclic aromatic ring.
106. A compound of claim 105, wherein said bicyclic aromatic ring is indenyl, quinoline or naphthyl.
107. A compound of claim 100, wherein Z is a bond, methyl, or ethyl.
108. A compound of claim 100, wherein n is 0.
109. A compound of claim 100, wherein X_1 and X_2 are both O.
110. A compound selected from the group consisting of
2-cyanoimino-3-ethyl-1-[1-(*p*-phenylbenzyl)-4-piperidiny]-1,3-dihydro-2H-benzimidazole;
2-cyanoimino-3-ethyl-1-[1-(*p*-benzyloxybenzyl)-4-piperidiny]-1,3-dihydro-2H-benzimidazole;
2-cyanoimino-3-ethyl-1-[1-(naphth-2-yl-methyl)-4-piperidiny]-1,3-dihydro-2H-benzimidazole;
2-cyanoimino-3-ethyl-1-[1-(4-propylcyclohexyl)-4-piperidiny]-1,3-dihydro-2H-benzimidazole;
2-cyanoimino-3-ethyl-1-[1-[4-(2-propyl)-cyclohexyl]-4-piperidiny]-1,3-dihydro-2H-benzimidazole;
2-cyanoimino-3-ethyl-1-[1-(decahydro-2-naphthyl)-4-piperidiny]-1,3-dihydro-2H-benzimidazole;
2-cyanoimino-3-ethyl-1-[1-(cyclooctyl)-4-piperidiny]-1,3-dihydro-2H-benzimidazole;
2-cyanoimino-3-ethyl-1-[1-(10,11-dihydro-5H-dibenzo[a,d]-cyclohepten-5-yl)-4-piperidiny]-1,3-dihydro-2H-benzimidazole;
2-cyanoimino-3-ethyl-1-[1-(3,3-Bis(phenyl)propyl)-4-piperidiny]-1,3-dihydro-2H-benzimidazole;
2-cyanoimino-3-ethyl-1-[1-(1,2,3,4-tetrahydronaphthyl)-4-piperidiny]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(5-methylhex-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(norbornan-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(1,3-dihydroinden-2-yl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-ethyl-1-[1-(cyclooctylmethyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole; and

pharmaceutically acceptable salts thereof.

111. A compound selected from the group consisting of

2-cyanoimino-3-(2-hydroxy)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-methoxycarbonylmethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-cyanomethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-butyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-(2-methanesulfonamido)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-acetomido-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-carboxymethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-3-(2-dimethylamino)ethyl-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;

2-cyanoimino-1-[1-(cyclooctyl)-3-hydroxymethyl-4-piperidinyl]-1,3-dihydro-2H-benzimidazole;


2-cyanoimino-1-[1-(cyclooctyl)-4-piperidinyl]-1,3-dihydro-2H-7-azabenzimidazole;

2-cyanoimino-1-[1-(cyclooctyl)-2,6-ethano-4-one-4-piperidinyl]-1,3-dihydro-2H-benzimidazole; and

pharmaceutically acceptable salts thereof and solvates thereof.

112. A pharmaceutical composition comprising a compound of claim 94 and at least one pharmaceutically acceptable excipient.

113. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 94.
114. A method of modulating a pharmacological response from the ORL1 receptor comprising administering to a patient in need thereof an effective amount of a compound according to claim 94.
115. A pharmaceutical composition comprising a compound of claim 100 and at least one pharmaceutically acceptable excipient.
116. A method of treating pain comprising administering to a patient in need thereof, an effective amount of an analgesic compound according to claim 100.
117. A method of modulating a pharmacological response from the ORL1 receptor comprising administering an effective amount of a compound according to claim 100.
118. A compound of the formula (IVA):



(IVA)


wherein

n is an integer from 0 to 3;

R is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, and C₃₋₁₂ cycloalkyl;


R₂ is selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl and halogen, said alkyl optionally substituted with an oxo group;

ZR₁ is the following



wherein

Y₁ is R₃-(C₁-C₁₂)alkyl, R₄-aryl, R₅-heteroaryl, R₆-(C₃-C₁₂)cyclo-alkyl, R₇-(C₃-C₇)heterocycloalkyl, -CO₂(C₁-C₆)alkyl, CN or -C(O)NR₈R₉; Y₂ is hydrogen or Y₁; Y₃ is hydrogen or (C₁-C₆)alkyl; or Y₁, Y₂ and Y₃, together with the carbon to which they are attached, form one of the following structures:



wherein r is 0 to 3; w and u are each 0-3, provided that the sum of w and u is 1-3; c and d are independently 1 or 2; s is 1 to 5; and ring E is a fused R_4 -phenyl or R_5 -heteroaryl ring;

R_{10} is 1 to 3 substituents independently selected from the group consisting of H, (C_1-C_6) alkyl, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-NR_8R_9$ and $-(C_1-C_6)$ alkyl- NR_8R_9 ;

R_{11} is 1 to 3 substituents independently selected from the group consisting of R_{10} , $-CF_3$, $-OCF_3$, NO_2 and halo, or R_{11} substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;


R_8 and R_9 are independently selected from the group consisting of hydrogen, (C_1-C_6) alkyl, (C_3-C_{12}) cycloalkyl, aryl and aryl (C_1-C_6) alkyl;

R_3 is 1 to 3 substituents independently selected from the group consisting of H, R_4 -

aryl, R_6 -(C_3 - C_{12})cycloalkyl, R_5 -heteroaryl, R_7 -(C_3 - C_7)heterocycloalkyl, $-NR_8R_9$, $-OR_{12}$ and $-S(O)_{0.2}R_{12}$;

R_6 is 1 to 3 substituents independently selected from the group consisting of H, (C_1 - C_6)alkyl, R_4 -aryl, $-NR_8R_9$, $-OR_{12}$ and $-SR_{12}$;

R_4 is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C_1 - C_6)alkyl, R_{13} -aryl, (C_3 - C_{12})cycloalkyl, $-CN$, $-CF_3$, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-OCF_3$, $-NR_8R_9$, $-(C_1-C_6)$ alkyl- NR_8R_9 , $-NHSO_2R_8$, $-SO_2N(R_{14})_2$, $-SO_2R_8$, $-SOR_8$, $-SR_8$, $-NO_2$, $-CONR_8R_9$, $-NR_9COR_8$, $-COR_8$, $-COCF_3$, $-OCOR_8$, $-OCO_2R_8$, $-COOR_8$, $-(C_1-C_6)$ alkyl- $NHCOOC(CH_3)_3$, $-(C_1-C_6)$ alkyl- $NHCOCF_3$, $-(C_1-C_6)$ alkyl- $NHSO_2$ -(C_1-C_6)alkyl, $-(C_1-C_6)$ alkyl- $NHCONH$ -(C_1-C_6)alkyl and



wherein f is 0 to 6; or R_4 substituents on adjacent ring carbon atoms may together form a methylenedioxy or ethylenedioxy ring;

R_5 is 1 to 3 substituents independently selected from the group consisting of hydrogen, halo, (C_1 - C_6)alkyl, R_{13} -aryl, (C_3 - C_{12})cycloalkyl, $-CN$, $-CF_3$, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-OCF_3$, $-NR_8R_9$, $-(C_1-C_6)$ alkyl- NR_8R_9 , $-NHSO_2R_8$, $-SO_2N(R_{14})_2$, $-NO_2$, $-CONR_8R_9$, $-NR_9COR_8$, $-COR_8$, $-OCOR_8$, $-OCO_2R_8$ and $-COOR_8$;

R_7 is H, (C_1 - C_6)alkyl, $-OR_8$, $-(C_1-C_6)$ alkyl- OR_8 , $-NR_8R_9$ or $-(C_1-C_6)$ alkyl- NR_8R_9 ;

R_{12} is H, (C_1 - C_6)alkyl, R_4 -aryl, $-(C_1-C_6)$ alkyl- OR_8 , $-(C_1-C_6)$ alkyl- NR_8R_9 , $-(C_1-C_6)$ alkyl- SR_8 , or aryl (C_1 - C_6)alkyl;

R_{13} is 1-3 substituents independently selected from the group consisting of H, (C_1 - C_6)alkyl, (C_1 - C_6)alkoxy and halo;

R_{14} is independently selected from the group consisting of H, (C_1 - C_6)alkyl and R_{13} - $C_6H_4-CH_2$ -

and a pharmaceutically acceptable salt thereof.

119. A pharmaceutical composition comprising a compound of claim 118 and at least one pharmaceutically acceptable excipient.

120. A method of treating pain comprising administering to a patient in need thereof, an

effective amount of an analgesic compound according to claim 118.

121. A method of modulating a pharmacological response from the ORL1 receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 118.

122. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 94.

123. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 100.

124. A method of modulating a pharmacological response from an opioid receptor comprising administering to a patient in need thereof, an effective amount of a compound according to claim 118.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/12351

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/278, 300, 302, 307, 321, 322, 323, 395; 546/17, 116, 139, 198, 199, 201; 548/306.4

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS—structure

EAST/WEST—subclass and pain, opio?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database on STN CASDATA (Columbus, Ohio, USA) CA accession: no. 129:54361, HUFF et al. 'Preparation of benzisothiazolones and analogs as alpha.1C. adrenergic receptor antagonist' US pat. no. 5,760,054, (1998) see registry numbered compounds.	1, 2, 17
X ----- Y	Database on STN CASDATA (Columbus, Ohio, USA) Ca, accession no. 130:95479, SASAKI et al 'preparation of piperidine derivatives as cell adhesion inhibitors' JP 10330377 (1998) see registry numbered compounds	1, 6, 10, 11, 13, 23, 24 ----- 1-10, 12, 14-22, 24



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A"	document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

11 JUNE 2002

Date of mailing of the international search report

17 JUL 2002

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CELIA CHANG

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US02/12351

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---- Y	Database STN CASDATA, (Columbus, Ohio, USA) CA accession 124:176079, Huff et al. 'preparation of heterocycles as alpha-1c adrenergic receptor antagonists' WO 9528397 (1995) see registry numbered compounds.	1, 6, 10, 11, 13, 23, 24 ----- 1-10, 12, 14-22
X ---- Y	US 3,325,499 a (POOS et al.) 13 June 1967, see entire article especially claims 1-4.	30, 31, 36, 38, 42, 45, 46, 56-62 ----- 32-35, 37, 39-41, 43, 44, 47-55
X ---- Y	Database on STN CASDATA (Columbus, Ohio, USA) CA accession 81:77786, Klein et al. 'Potential analgesics' Arch. Pharm. (1974) v. 307 (5) pages 360-5, see registry numbered compounds.	30, 36-39, 42, 45, 46, 50-55 ----- 31- 35, 40, 41, 43-44, 47-48
X ---- Y	Database on STN CASDATA (Columbus, Ohio, USA) CA Accession no. 136:53656 Forbes 'A short and efficient synthesis of n-substituted indol-2-ones' Tetrahedron Let. (2001) v.42(39)pages 6943-6945, see registry numbered compounds.	30, 36-38, 42, 45- 46, 50-55 ----- 31-35, 39-41, 47- 49
X ---- Y	Database on STN CASDATA (Columbus, Ohio, USA) CA accession no. 135:195564, TSUSHIMA et al. 'preparation of phenoxyalkylamine derivative useful as opioid delta receptor agonists' WO 2001/60796 (2001) see registry numbered compounds.	30, 36, 38, 42, 45, 46, 50-55 ----- 31-35, 37, 39-41, 43, 44
P, X ----- P, Y	Database on STN CASDATA, (Columbus, Ohio, USA) CA accession no. 135:272955, TSUSHIMA et al. 'Preparation of diphenylalkylpiperidine derivatives useful as opioid delta receptor agonists' WO 2001/70689 (2001), see registry numbered compounds.	30, 36-38, 42, 45, 46, 50-55 ----- 31-35, 39-41, 43, 44, 56-62
Y	Database on STN CASDATA, (Columbus, Ohio, USA) CA accession no. 130:223167, BUDHU et al. 'preparation of piperidinylpyrrolidins as modulators of chemokine receptor' WO 99/09984 (1999) see registry numbered compounds.	30, 36
X ---- Y	Database on STN CASDATA (Columbus, Ohio, USA) CA accession no. 95:203828, BIANCHI et al. 'Compounds with antiulcer and antisecretory activity.' Eur. J. Med. Chem. (1981) 1694, pages 321-6, see registry numbered compounds.	63, 64, 69, 73, 76, 81, 87, 88 ----- 65-68, 70-75, 77- 80

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US02/12351

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---- Y	Database on STN CASDATA (Columbus, Ohio, USA) CA Accession no. 73:445405, OGATA et al. 'Organic photochemical reactions' Chem. Pharm Bull (1970) 18(5) pages 964-9, see registry numbered compounds.	63, 64, 69, 73, 76, 81
X	Database on STN CASDATA (Columbus, Ohio, USA) CA accession no. 95:177993, GUILLIANO et al. 'Spectroscopic studies of 2-oxo-2,3-dihydrobenzimidazole and some n-monosubstituted or disubstituted derivatives' App. Spectrosc. (1981) 3595) pages 486-8, see registry numbered compound.	63, 64, 69, 73, 76, 81
X ---- Y	Database on STN CASDATA (Columbus, Ohio, USA) CA accession no. 134:353434, NAKAMURA et al. 'synthesis of a regio-isomer of kealiiquinon' J. Chem. soc. japan (2001) (4) pages 429-436, see registry numbered compound.	63, 64, 69, 73, 76, 81, 87, 88 ----- 65-69, 70-72, 74, 75, 79, 80
Y	Database on STN CASDATA (Columbus, Ohio, USA) CA accession no. 125:58298, SUZUKI et al. 'preparation of heterocycle containing amides as 5HT4 agonists' WO 96/05166, (1996) see registry numbered compounds	69, 81
Y	US 5,574,044 A (THOMPSON et al.) 12 November 1996, see entire document especially col. 20, first three compounds.	63, 69, 81, 87, 88
X ---- Y	US 4,215,119 A (MENTRUP et al.) 29 July 1980, see entire article especially columns 78-82, compounds.	63, 69, 81, 87, 88 ----- 64-68, 80,
X ---- Y	US 5,661,169 A (DIMALTA et al.) 26 August 1997, see entire article especially compounds at columns 19-20.	63, 69, 87, ----- ----- 64-68, 70-80
X ---- Y	US 4,410,528 A (TERANISHI ET AL. et al.) 18 October 1983, see entire article especially column 39-40 examples 11-12.	94, 100, 107, 108, 118 ----- 95-99, 101, 103-111

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/12351

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/12351

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

A61K 31/44, 31/48, 31/415, 31/445; C07D 211/04, 211/06, 235/24, 401/04, 401/06, 403/04, 403/06, 471/02, 498/02

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

514/278, 300, 302, 307, 321, 322, 323, 395; 546/17, 116, 139, 198, 199, 201; 548/306.4

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-29, drawn to benzoxazolonyl compounds composition and method of use.

Group II, claims 30-62, drawn to indolonyl compounds composition and method of use.

Group III, claims 63-93, drawn to benzimidazolonyl compounds composition and method of use.

Group IV, claims 94-124, drawn to benzimidazocyanimidyl compounds composition and method of use.

The inventions listed as Groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: each group of compounds have different structural cores that do not share similarity with each other.